TUHEP-TH-07157

arXiv:0705.0115v2 [hep-ph] 2 May 2007

Dynamical Computation on Coefficients of Electroweak Chiral Lagrangian from One-doublet and Topcolor-assisted Technicolor Models

Hong-Hao Zhang*, Shao-Zhou Jiang†, and Qing Wang‡§

Department of Physics, Tsinghua University, Beijing 100084, China¶

Center for High Energy Physics, Tsinghua University, Beijing 100084, China

Abstract

Based on previous studies deriving the chiral Lagrangian for pseudo scalar mesons from the first principle of QCD, we derive the electroweak chiral Lagrangian and build up a formulation for computing its coefficients from one-doublet technicolor model and a schematic topcolor-assisted technicolor model. We find that the coefficients of the electroweak chiral Lagrangian for the topcolor-assisted technicolor model are divided into three parts: direct TC2 interaction part, TC1 and TC2 induced effective Z' particle contribution part, and ordinary quarks contribution part. The first two parts are computed in this paper and we show that the direct TC2 interaction part is the same as that in the one-doublet technicolor model, while effective Z' contributions are at least proportional to the p^2 order parameter β_1 in the electroweak chiral Lagrangian and typical features of topcolor-assisted technicolor model are that it only allows positive T and U parameters and the T parameter varies in the range $0 \sim 1/(25\alpha)$, the upper bound of T parameter will decrease as long as Z' mass become large. The S parameter can be either positive or negative depending on whether the Z' mass is large or small. The Z' mass is also bounded above and the upper bound depend on value of T parameter. We obtain the values for all the coefficients of the electroweak chiral Lagrangian up to order of p^4.

PACS numbers: 11.10.Lm, 11.30.Rd, 12.10.Dm, 12.60.Nz

---

* Email: zhanghonghao@tsinghua.org.cn.

† Email: jsz@mails.tsinghua.edu.cn.

‡ Email: wangq@mail.tsinghua.edu.cn.

§ corresponding author

¶ mailing address

Contents

I. Introduction 3

II. Derivation of the Electroweak Chiral Lagrangian from the One-doublet Technicolor Model 5

A. The Gasser-Leutwyler's Prescription 7

B. The Dynamical Computation Prescription 8

1. Schwinger-Dyson Equation for Techniquark Propagator 11

2. Effective Action 15

C. Comparison and Discussion on Two Prescriptions 19

III. Derivation of the Electroweak Chiral Lagrangian from a Topcolor-assisted Technicolor Model 20

A. Topcolor Symmetry Breaking: the Contribution of $SU(3)_{\rm TC1}$ 22

1. The Gap Equation 25

2. Low Energy Expansion for $iS_{\rm TC1}[0, B^A_\mu, B_\mu, Z'_\mu]$ 28

B. Electroweak Symmetry Breaking: the Contribution of $SU(3)_{\rm TC2}$ 30

C. Integrate out of Colorons 35

D. Integrate out of Z' 37

IV. Conclusion 43

Acknowledgments 45

A. Two Equivalent EWCL Formalisms 46

B. $S^{(4c,B^A)}_{\rm TC1}[B^A]$, $S^{(4c,B)}_{\rm TC1}[B]$, $S^{(4c,Z')}_{\rm TC1}$, $S^{(4c,B^AZ')}_{\rm TC1}[B^A, Z']$, $S^{(4c,BZ')}_{\rm TC1}[B, Z']$ 48

C. $S^0_{\rm coloron}[B^A, Z']$ **and** $S^{\rm int}_{\rm coloron}[B^A, Z']$ 50

References 51

I. INTRODUCTION

The electroweak symmetry breaking mechanism (EWSBM) remains an intriguing puzzle for particle physics, although the Standard Model (SM) provides us with a version of it through introducing a higgs boson into the theory which suffers from triviality and unnaturalness problems. Beyond SM, numerous new physics models are invented which exhibit many alternative EWSBMs. With the present situation that higgs is still not found in experiment, all new physics models at low energy region should be described by a theory which not only must match all present experiment data, but also have no higgs. This theory is the well-known electroweak chiral Lagrangian (EWCL) [1, 2, 3] which offers the most general and economic description of electroweak interaction at low energy region. With EWCL, new physics models at low energies can be parameterized by a set of coefficients, it universally describes all possible electroweak interactions among existing particles and offers a model independent platform for us to investigate various kinds EWSBMs. Starting from this platform, further phenomenological research focus on finding effective physical processes to fix the certain coefficients of EWCL [4, 5, 6], and theoretical studies concentrate on consistency of EWCL itself such as gauge invariance [7] and computing the values of the coefficients for SM with heavy higgs [8]. Systematic theoretical computation of the EWCL coefficients for other new physics models have not been presented in the literature. The possible reasons are that for weakly coupled models, since one can perform perturbative computations, people prefer to directly discuss physics from the model and then are reluctant to pay the extra price to compute EWCL coefficients. While for strongly coupled models, non-perturbative difficulties for a long time prevent people to perform dynamical computations, only for special coefficients such as S parameter, some non-perturbative technique may be applied to perform calculations [9] or for special QCD-like technicolor models, in terms of their similarities with QCD, one can estimate the coefficients of EWCL in terms of their partners fixed by experimental data in corresponding QCD chiral Lagrangian. The estimation of EWCL coefficients for various models is of special importance in the sense that at present we already have some quantitative constraints on them, such as those for the S, T, U parameters and more generally for anomalous triple and quartic couplings [10], along with the experimental progress, more constraints will be obtained. Once we know the values of the coefficients for detailed models, these constraints can directly be used to judge the correctness of the model. It is the purpose of this paper to develop a formulation to systematically compute coefficients of EWCL for strongly coupled new physics models. For simplicity, in this work we only discuss the bosonic part of EWCL and leave the matter part for future investigations. The basis of our formulation is the knowledge and experiences we obtained previously from a series of works deriving the chiral Lagrangian for pseudo scalar mesons from QCD first principles [11] and calculate coefficients in it [12, 13, 14], where we found confidence and reliability in this work. In fact, the formal derivation from a general underlying technicolor model to EWCL was already achieved in Ref.[15] in which except deriving EWCL, coefficients of EWCL are formally expressed in terms of Green's functions in underlying technicolor model. Once we know how to compute these Green's functions, we obtain the corresponding EWCL coefficients. The pity is that the computation is nonperturbative, therefore not easy to achieve, it is the aim of this paper to solve this nonperturbative dynamical computation problems.

As the first step of performing dynamical computations, we especially care about the reliability of the formulation we will develop. We take one-doublet technicolor model [16, 17, 18, 19] as the prototype to build up our formulation. Although this model as the earliest and simplest dynamical electroweak symmetry breaking model was already denied by experiment in the sense that it results in too large a value for the S parameter, but due to the following reasons, we still start our investigations from here. First, it is similar as QCD in the theory structures which enable us to easily generalize the techniques developed in dealing with the QCD chiral Lagrangian to this model and we call this generalized formulation the dynamical computation prescription. Second due to their similarities with conventional QCD, the coefficients of their EWCLs can be estimated by just scaling-up corresponding coefficients in QCD Gasser-Leutwyler chiral Lagrangian for pseudo scalar mesons [20] and we call this formulation the Gasser-Leutwyler's prescription which naively is only applicable for those QCD-like models. So for QCD-like models, we have two prescriptions which enable us to compare them with each other to check the correctness and increase the reliability of our formulations. Beyond the traditional one-doublet technicolor model, we choose the topcolor-assisted technicolor model as the first real practice model to perform our computations. The reason to take it is that this model is not QCD-like and is active on the market now which does not seriously contradict with experimental data as the case of one-doublet technicolor model and the dynamics responsible for electroweak symmetry breaking is similar to that in the one-doublet technicolor model. We will find that the coefficients for topcolor-assisted technicolor model can be divided into three parts: direct TC2 interaction part,TC1 and TC2 induced effective Z' particle contribution part and ordinary quarks contribution part. The first two parts are computed in this paper and we show that direct TC2 interaction part is same as that in one-doublet technicolor model, while TC1 and TC2 induced effective Z' particle contributions is at least proportional to p^2 order parameter β_1 in EWCL and typical features of topcolor-assisted technicolor model are that it only allows positive T and U parameters and the T parameter varies in the range $0 \sim 1/(25\alpha)$,the upper bound of T parameter will decrease as long as Z' mass become large. The S parameter can be either positive or negative depending on whether the Z' mass is large or small. The Z' mass is also bounded above and the upper bound depend on value of T parameter. We obtain the values for all the coefficients of the electroweak chiral Lagrangian up to order of p^4.

This paper is organized as follows. Section II is the basics of the work in which we discuss one-doublet technicolor model. We first review the Gasser-Leutwyler's prescription, then build up our dynamical computation prescription, we show how to consistently set in the dynamical computation equation (the Schwinger-Dyson equation) into our formulation. We make comparison between two prescriptions to check validity of the results from our dynamical computation prescription. Section III is the main part of this work in which we apply our formulations developed in one-doublet technicolor model to topcolor-assisted technicolor model. We perform dynamical calculations on technicolor interactions and then integrate out colorons and Z' to compute EWCL coefficients. Since this is the first time to systematically perform dynamical computations on the strongly coupled models, we emphasize the technical side more than physics analysis and display the computation procedure a little bit more in detail. Section IV is the conclusion. In the appendices, we list some requisite formulae.

II. DERIVATION OF THE ELECTROWEAK CHIRAL LAGRANGIAN FROM THE ONE-DOUBLET TECHNICOLOR MODEL

Consider the one-doublet technicolor (TC) model proposed by Weinberg and Susskind independently [16, 17, 18, 19]. The techniquarks are assigned to $(SU(N)_{TC}, SU(3)_C, SU(2)_L, U(1)_Y)$ as $\psi_L \sim (N,1,2,0)$, $U_R = (1/2 + \tau^3/2)\psi_R \sim$

$(N,1,1,1/2)$, $D_R=(1/2-\tau^3/2)$, $\psi_R\sim(N,1,1,-1/2)$. With these assignments, the techniquarks have electric charges as defined by $Q=T_3+Y$, of $+1/2$ for U and $-1/2$ for D. It can be shown below, by dynamical analysis through the Schwinger-Dyson equation, that the $SU(N)_{TC}$ interaction induces the techniquark condensate $\langle\bar{\psi}\psi\rangle\neq 0$, which will trigger the electroweak symmetry breaking $SU(2)_L\times U(1)_Y\to U(1)_{\rm EM}$. Neglecting ordinary fermions and gluons, we focus on the action of the techniquark, technicolor-gauge-boson and electroweak-gauge-boson sector, *i.e.* the electroweak symmetry breaking sector (SBS) of this model,

$$
\begin{aligned}
S_{\rm SBS} \;=\; \int d^4x\Big[&-\frac{1}{4}F^\alpha_{\mu\nu}F^{\alpha,\mu\nu}-\frac{1}{4}W^a_{\mu\nu}W^{a,\mu\nu}-\frac{1}{4}B_{\mu\nu}B^{\mu\nu}\\
&+\bar{\psi}\Big(i\not\partial-g_{\rm TC}t^\alpha\not{G}^\alpha-g_2\frac{\tau^a}{2}\not{W}^aP_L-g_1\frac{\tau^3}{2}\not{B}P_R\Big)\psi\Big]\,, \qquad (1)
\end{aligned}
$$

where $g_{\rm TC}$, g_2 and g_1 (G^α_μ, W^a_μ and B_μ) are the coupling constants (gauge fields) of $SU(N)_{TC}\times SU(2)_L\times U(1)_Y$ with technicolor index α ($\alpha=1,2,\ldots,N^2-1$) and weak index a ($a=1,2,3$) respectively; and $F^\alpha_{\mu\nu}$, $W^a_{\mu\nu}$ and $B_{\mu\nu}$ are the corresponding field strength tensors; t^α ($\alpha=1,2,\ldots,N^2-1$) are the generators for the fundamental representation of $SU(N)_{TC}$, while τ^a ($a=1,2,3$) are Pauli matrices; and the left and right chirality projection operators $P_{\substack{L\\R}}=(1\mp\gamma_5)/2$.

To derive low energy effective electroweak chiral Lagrangian from the one-doublet TC model, we need to integrate out the technigluons and techniquarks above the electroweak scale which can be formulated as

$$
\int\mathcal{D}G^\alpha_\mu\mathcal{D}\bar{\psi}\mathcal{D}\psi\exp\bigg(iS_{\rm SBS}\big[G^\alpha_\mu,W^a_\mu,B_\mu,\bar{\psi},\psi\big]\bigg)=\int\mathcal{D}\mu(U)\exp\bigg(iS_{\rm eff}[U,W^a_\mu,B_\mu]\bigg)\,, \qquad (2)
$$

where $U(x)$ is a dimensionless unitary unimodular matrix field in the electroweak chiral Lagrangian, and $\mathcal{D}\mu(U)$ denotes the corresponding functional integration measure.

As mentioned in previous section, there are two different approaches, one is the Gasser-Leutwyler's prescription, the other is the dynamical computation prescription. The second approach we developed in this paper is relatively easy to be generalized to more complicated theories. We will compare the results obtained in both approaches.

A. The Gasser-Leutwyler's Prescription

As we will see, it is easy to relate QCD-like models to chiral Lagrangian using the Gasser-Leutwyler's prescription. To begin, we substitute (1) into the left-hand side of Eq.(2) and the result path integral involved technicolor interaction is analogous to QCD and then we can use technique developed by Gasser and Leutwyler relating it with the path integral of chiral Lagrangian for goldstone bosons induced from SBS [20],

$$\frac{\int\mathcal{D}G^{\alpha}_{\mu}\mathcal{D}\bar{\psi}\mathcal{D}\psi\exp\left\{i\int d^4x\left[-\frac{1}{4}F^{\alpha}_{\mu\nu}F^{\alpha,\mu\nu}+\bar{\psi}\left(i\not{\partial}-g_{\rm TC}t^{\alpha}\not{G}^{\alpha}-g_2\frac{\tau^a}{2}\not{W}^aP_L-g_1\frac{\tau^3}{2}\not{B}P_R\right)\psi\right]\right\}}{\int\mathcal{D}\bar{\psi}\mathcal{D}\psi\exp\{i\int d^4x\bar{\psi}[i\not{\partial}-g_2\frac{\tau^a}{2}\not{W}^aP_L-g_1\frac{\tau^3}{2}\not{B}P_R]\psi\}}$$
$$=\int\mathcal{D}\mu(\tilde{U})\exp\{iS_{\text{TC-induced eff}}[\tilde{U},W,B]\}\ , \qquad (3)$$

in which the denominator of the left hand side of above equation is introduced to insure the technicolor-induced chiral effective action $S_{\text{TC-induced eff}}[\tilde{U},W,B]$ normalized as zero when we switch off technicolor interactions by setting $g_{\rm TC}=0$ and $S_{\text{TC-induced eff}}[\tilde{U},W,B]$ can be written as

$$\begin{aligned}S_{\text{TC-induced eff}}[\tilde{U},W,B] \;=\; &\int d^4x\bigg[\frac{(F_0^{\rm 1D})^2}{4}{\rm tr}[(\nabla^{\mu}\tilde{U}^{\dagger})(\nabla_{\mu}\tilde{U})]+L_1^{\rm 1D}[{\rm tr}(\nabla^{\mu}\tilde{U}^{\dagger}\nabla_{\mu}\tilde{U})]^2\\&+L_2^{\rm 1D}{\rm tr}[\nabla_{\mu}\tilde{U}^{\dagger}\nabla_{\nu}\tilde{U}]{\rm tr}[\nabla^{\mu}\tilde{U}^{\dagger}\nabla^{\nu}\tilde{U}]+L_3^{\rm 1D}{\rm tr}[(\nabla^{\mu}\tilde{U}^{\dagger}\nabla_{\mu}\tilde{U})^2]\\&-iL_9^{\rm 1D}{\rm tr}[F^R_{\mu\nu}\nabla^{\mu}\tilde{U}\nabla^{\nu}\tilde{U}^{\dagger}+F^L_{\mu\nu}\nabla^{\mu}\tilde{U}^{\dagger}\nabla^{\nu}\tilde{U}]+L_{10}^{\rm 1D}{\rm tr}[\tilde{U}^{\dagger}F^R_{\mu\nu}\tilde{U}F^{L,\mu\nu}]\\&+H_1^{\rm 1D}{\rm tr}[F^R_{\mu\nu}F^{R,\mu\nu}+F^L_{\mu\nu}F^{L,\mu\nu}]\bigg]+O(p^6)\ ,\end{aligned}\qquad (4)$$

where the coefficients $F_0^{\rm 1D}$, $L_1^{\rm 1D}$, $L_2^{\rm 1D}$, $L_3^{\rm 1D}$, $L_{10}^{\rm 1D}$, $H_1^{\rm 1D}$ arise from $SU(N)_{\rm TC}$ dynamics at the scale of 250 GeV, and where

$$\begin{aligned}&\nabla_{\mu}\tilde{U}\equiv\partial_{\mu}\tilde{U}-ir_{\mu}\tilde{U}+i\tilde{U}l_{\mu}\ ,\qquad \nabla_{\mu}\tilde{U}^{\dagger}=-\tilde{U}^{\dagger}(\nabla_{\mu}\tilde{U})\tilde{U}^{\dagger}=\partial_{\mu}\tilde{U}^{\dagger}-il_{\mu}\tilde{U}^{\dagger}+i\tilde{U}^{\dagger}r_{\mu}\ ,\\&F^R_{\mu\nu}\equiv i[\partial_{\mu}-ir_{\mu},\partial_{\nu}-ir_{\nu}]\quad F^L_{\mu\nu}\equiv i[\partial_{\mu}-il_{\mu},\partial_{\nu}-il_{\nu}]\quad r_{\mu}\equiv-g_1\frac{\tau^3}{2}B_{\mu}\quad l_{\mu}\equiv-g_2\frac{\tau^a}{2}W^a_{\mu}\ .\end{aligned}\qquad (5)$$

Note that conventional $\tilde{U}$ field in Eq.(4) given in second paper of [20] is 3×3 unitary matrix. However, for the $SU(2)_L\times SU(2)_R$ electroweak chiral Lagrangian we considered in this paper, $\tilde{U}$ is a 2×2 unitary matrix, and thus the $L_1^{\rm 1D}$ term and the $L_3^{\rm 1D}$ term in present situation are linearly related,

$$L_3^{\rm 1D}{\rm tr}[(\nabla^{\mu}\tilde{U}^{\dagger}\nabla_{\mu}\tilde{U})^2]=L_3^{\rm 1D}{\rm tr}\Big\{\big[\tilde{U}^{\dagger}(\nabla^{\mu}\tilde{U})\tilde{U}^{\dagger}(\nabla^{\mu}\tilde{U})\big]^2\Big\}=\frac{L_3^{\rm 1D}}{2}\Big\{{\rm tr}\big[\tilde{U}^{\dagger}(\nabla^{\mu}\tilde{U})\tilde{U}^{\dagger}(\nabla^{\mu}\tilde{U})\big]\Big\}^2\quad(6)$$

Comparing covariant derivative for $\tilde{U}$ given in (5) and covariant derivative given in Ref.[2], we find we must recognize $\tilde{U}^\dagger = U$, $\nabla_\mu \tilde{U}^\dagger = D_\mu U$, $F^R_{\mu\nu} = -g_1 \frac{\tau^3}{2} B_{\mu\nu}$ and $F^L_{\mu\nu} = -g_2 \frac{\tau^a}{2} W^a_{\mu\nu}$. Substitute them back into Eq.(4), we obtain

$$\begin{aligned} S_{\text{TC-induced eff}}[U,W,B] \;=\; \int d^4x \Big[& -\frac{(F_0^{\rm 1D})^2}{4} \mathrm{tr}(X_\mu X^\mu) + (L_1^{\rm 1D} + \frac{L_3^{\rm 1D}}{2})[\mathrm{tr}(X_\mu X^\mu)]^2 \\ & + L_2^{\rm 1D}[\mathrm{tr}(X_\mu X_\nu)]^2 - i\frac{L_9^{\rm 1D}}{2} g_1 B_{\mu\nu} \mathrm{tr}(\tau^3 X^\mu X^\nu) - iL_9^{\rm 1D} \mathrm{tr}(\overline{W}_{\mu\nu} X^\mu X^\nu) \\ & + \frac{L_{10}^{\rm 1D}}{2} g_1 B_{\mu\nu} \mathrm{tr}(\tau^3 \overline{W}^{\mu\nu}) + \frac{H_1^{\rm 1D}}{2} g_1^2 B_{\mu\nu} B^{\mu\nu} + H_1^{\rm 1D} \mathrm{tr}(\overline{W}_{\mu\nu} \overline{W}^{\mu\nu}) \Big] \,. \end{aligned} \tag{7}$$

where

$$X_\mu \equiv U^\dagger (D_\mu U) \qquad\qquad \overline{W}_{\mu\nu} \equiv U^\dagger g_2 W_{\mu\nu} U \;, \tag{8}$$

We have reformulated the EWCL in terms of X_μ and τ^3 instead of V_μ and T in Ref.[2], the corresponding relations are given in Appendix A. Comparing (7) with the standard electroweak chiral Lagrangian given in Ref.[2], we find

$$\begin{aligned} & f^2 = (F_0^{\rm 1D})^2 \,, \qquad \beta_1 = 0 \,, \qquad \alpha_1 = L_{10}^{\rm 1D} \,, \qquad \alpha_2 = \alpha_3 = -\frac{L_9^{\rm 1D}}{2} \,, \qquad \alpha_4 = L_2^{\rm 1D} \,, \\ & \alpha_5 = L_1^{\rm 1D} + \frac{L_3^{\rm 1D}}{2} \,, \qquad \alpha_i = 0 \;\; (i = 6, 7, \ldots, 14) \,. \end{aligned} \tag{9}$$

Note that in (7), the term with coefficient $H_1^{\rm 1D}$ do not affect the result α_i coefficients. Not like the original case of QCD, $H_1^{\rm 1D}$ now is a finite constant. The divergences are from the term $\mathrm{Tr}\log(i\not\partial - g_2 \frac{\tau^a}{2} \not{W}^a P_L - g_1 \frac{\tau^3}{2} \not{B} P_R)$ in (3) which will only contribute $\mathrm{tr}(W_{\mu\nu} W^{\mu\nu})$ and $B_\mu B^\mu$ terms with divergent coefficients due to gauge invariance. These divergent coefficients in combined with $H_1^{\rm 1D}$ will cause wave function renormalization corrections for W^a_μ and B_μ fields which further lead redefinitions of W^a_μ and B_μ fields and their gauge couplings g_2 and g_1. This renormalization procedure will have no effects on our EWCL, since all electroweak gauge fields appeared in EWCL are as form of $g_2 W^a_\mu$ and $g_1 B_\mu$ which are renormalization invariant quantities. Due to this consideration, in rest of this paper, we just left the wave function corrections to electroweak gauge fields W^a_μ and B_μ in the theory and skip the corresponding renormalization procedure.

B. The Dynamical Computation Prescription

Now we develop a dynamical computation program and we will apply it to more complicated model in next section.

We first review the derivation process given in Ref.[15] and start with introducing a local 2×2 operator $O(x)$ as $O(x) \equiv \mathrm{tr}_{lc}[\psi_L(x)\bar{\psi}_R(x)]$ with tr_{lc} is the trace with respect to Lorentz and technicolor indices. The transformation of $O(x)$ under $SU(2)_L \times U(1)_Y$ is $O(x) \to V_L(x)O(x)V_R^\dagger(x)$. Then we decompose $O(x)$ as $O(x) = \xi_L^\dagger(x)\sigma(x)\xi_R(x)$ with the $\sigma(x)$ represented by a hermitian matrix describes the modular degree of freedom; while $\xi_L(x)$ and $\xi_R(x)$ represented by unitary matrices describe the phase degree of freedom of $SU(2)_L$ and $U(1)_Y$ respectively. Their transformation under $SU(2)_L \times U(1)_Y$ are $\sigma(x) \to h(x)\sigma(x)h^\dagger(x)$, $\xi_L(x) \to h(x)\xi_L(x)V_L^\dagger(x)$ and $\xi_R(x) \to h(x)\xi_R(x)V_R^\dagger(x)$ with $h(x) = e^{i\theta_h(x)\tau^3/2}$ belongs to an induced hidden local $U(1)$ symmetry group. Now we define a new field $U(x)$ as $U(x) \equiv \xi_L^\dagger(x)\xi_R(x)$ which is the nonlinear realization of the goldstone boson fields in the electroweak chiral Lagrangian. Subtracting the $\sigma(x)$ field, we find that the present decomposition results in a constraint $\xi_L(x)O(x)\xi_R^\dagger(x) - \xi_R(x)O^\dagger(x)\xi_L^\dagger(x) = 0$, the functional expression of it is

$$\int \mathcal{D}\mu(U)\mathcal{F}[O]\delta(\xi_L O\xi_R^\dagger - \xi_R O^\dagger\xi_L^\dagger) = \text{const.}\,, \tag{10}$$

where $\mathcal{D}\mu(U)$ is an effective invariant integration measure; $\mathcal{F}[O]$ only depends on O, and it compensates the integration to make it to be a constant. It is easy to show that $\mathcal{F}[O]$ is invariant under $SU(2)_L \times U(1)_Y$ transformations. Substituting Eq.(10) into the left-hand side of Eq.(2), we have

$$\int \mathcal{D}G_\mu^\alpha \mathcal{D}\bar{\psi}\mathcal{D}\psi \exp\Big(iS_{\text{SBS}}[G_\mu^\alpha, W_\mu^a, B_\mu, \bar{\psi}, \psi]\Big) = \int \mathcal{D}\mu(U)\exp\Big(iS_{\text{eff}}[U, W_\mu^a, B_\mu]\Big)\,, \tag{11}$$

where

$$\begin{aligned}
&S_{\text{eff}}[U, W_\mu^a, B_\mu] \qquad\qquad (12)\\
&= \int d^4x\Big(-\frac{1}{4}W_{\mu\nu}^a W^{a,\mu\nu} - \frac{1}{4}B_{\mu\nu}B^{\mu\nu}\Big) - i\log\int \mathcal{D}G_\mu^\alpha \mathcal{D}\bar{\psi}\mathcal{D}\psi\,\mathcal{F}[O]\delta(\xi_L O\xi_R^\dagger - \xi_R O^\dagger\xi_L^\dagger)\\
&\quad \times \exp\bigg\{i\int d^4x\Big[-\frac{1}{4}F_{\mu\nu}^\alpha F^{\alpha,\mu\nu} + \bar{\psi}\Big(i\partial\!\!\!/ - g_{\text{TC}}t^\alpha G\!\!\!/^\alpha - g_2\frac{\tau^a}{2}W\!\!\!\!/^a P_L - g_1\frac{\tau^3}{2}B\!\!\!/P_R\Big)\psi\Big]\bigg\}
\end{aligned}$$

To match the correct normalization, we introduce in the argument of logarithm function the normalization factor $\int \mathcal{D}\bar{\psi}\mathcal{D}\psi e^{i\int d^4x\bar{\psi}(i\partial\!\!\!/ - g_2\frac{\tau^a}{2}W\!\!\!\!/^a P_L - g_1\frac{\tau^3}{2}B\!\!\!/P_R)\psi} = \exp\mathrm{Tr}\log(i\partial\!\!\!/ - g_2\frac{\tau^a}{2}W\!\!\!\!/^a P_L - g_1\frac{\tau^3}{2}B\!\!\!/P_R)$ and then take a special $SU(2)_L \times U(1)_Y$ rotation, as $V_L(x) = \xi_L(x)$ and $V_R(x) =$

$\xi_R(x)$, on both numerator and denominator of the normalization factor

$$\begin{aligned}
&S_{\rm eff}[U,W_\mu^a,B_\mu] \qquad\qquad (13)\\
&= \int d^4x(-\frac{1}{4}W^a_{\mu\nu}W^{a,\mu\nu}-\frac{1}{4}B_{\mu\nu}B^{\mu\nu})+\mathrm{Tr}\log(i\partial\!\!\!/-g_2\frac{\tau^a}{2}W\!\!\!\!/^{\,a}P_L-g_1\frac{\tau^3}{2}B\!\!\!/P_R)\\
&-i\log\frac{\int\mathcal{D}G^\alpha_\mu\mathcal{D}\bar{\psi}_\xi\mathcal{D}\psi_\xi\,\mathcal{F}[O_\xi]\delta(O_\xi-O^\dagger_\xi)e^{i\int d^4x[-\frac{1}{4}F^\alpha_{\mu\nu}F^{\alpha,\mu\nu}+\bar{\psi}_\xi(i\partial\!\!\!/-g_{\rm TC}t^\alpha G\!\!\!/^\alpha-g_2\frac{\tau^a}{2}W\!\!\!\!/^{\,a}_\xi P_L-g_1\frac{\tau^3}{2}B\!\!\!/_\xi P_R)\psi_\xi]}}{\int\mathcal{D}\bar{\psi}_\xi\mathcal{D}\psi_\xi e^{i\int d^4x\bar{\psi}_\xi(i\partial\!\!\!/-g_2\frac{\tau^a}{2}W\!\!\!\!/^{\,a}_\xi P_L-g_1\frac{\tau^3}{2}B\!\!\!/_\xi P_R)\psi_\xi}}\,,
\end{aligned}$$

where rotated fields are denoted as follows

$$\begin{aligned}
&\psi_\xi=P_L\xi_L(x)\psi_L(x)+P_R\xi_R(x)\psi_R(x)\,,\qquad O_\xi(x)\equiv\xi_L(x)O(x)\xi^\dagger_R(x)\,, \qquad (14)\\
&g_2\frac{\tau^a}{2}W^a_{\xi,\mu}(x)\equiv\xi_L(x)[g_2\frac{\tau^a}{2}W^a_\mu(x)-i\partial_\mu]\xi^\dagger_L(x)\quad g_1\frac{\tau^3}{2}B_{\xi,\mu}(x)\equiv\xi_R(x)[g_1\frac{\tau^3}{2}B_\mu(x)-i\partial_\mu]\xi^\dagger_R(x)\,.
\end{aligned}$$

In (13), the possible anomalies caused by this special chiral rotation are canceled between the numerator and the denominator. Thus Eq.(13) can be written as

$$S_{\rm eff}[U,W^a_\mu,B_\mu]=\int d^4x(-\frac{1}{4}W^a_{\mu\nu}W^{a,\mu\nu}-\frac{1}{4}B_{\mu\nu}B^{\mu\nu})+S_{\rm anom}[U,W^a_\mu,B_\mu]+S_{\rm norm}[U,W^a_\mu,B_\mu],\quad(15a)$$

where

$$\begin{aligned}
S_{\rm norm}[U,W^a_\mu,B_\mu] \;=\; &-i\log\int\mathcal{D}G^\alpha_\mu\mathcal{D}\bar{\psi}_\xi\mathcal{D}\psi_\xi\,\mathcal{F}[O_\xi]\delta(O_\xi-O^\dagger_\xi)\exp\bigg\{i\int d^4x\bigg[-\frac{1}{4}F^\alpha_{\mu\nu}F^{\alpha,\mu\nu}\\
&+\bar{\psi}_\xi\bigg(i\partial\!\!\!/-g_{\rm TC}t^\alpha G\!\!\!/^\alpha-g_2\frac{\tau^a}{2}W\!\!\!\!/^{\,a}_\xi P_L-g_1\frac{\tau^3}{2}B\!\!\!/_\xi P_R\bigg)\psi_\xi\bigg]\bigg\}\,. \qquad (15b)
\end{aligned}$$

and

$$\begin{aligned}
iS_{\rm anom}[U,W^a_\mu,B_\mu] \;=\; &\mathrm{Tr}\log(i\partial\!\!\!/-g_2\frac{\tau^a}{2}W\!\!\!\!/^{\,a}P_L-g_1\frac{\tau^3}{2}B\!\!\!/P_R)\\
&-\mathrm{Tr}\log(i\partial\!\!\!/-g_2\frac{\tau^a}{2}W\!\!\!\!/^{\,a}_\xi P_L-g_1\frac{\tau^3}{2}B\!\!\!/_\xi P_R)\,, \qquad (15c)
\end{aligned}$$

It is worthwhile to show the transformations of the rotated fields under $SU(2)_L\times U(1)_Y$ are $\psi_\xi(x)\to h(x)\psi_\xi(x)$, $O_\xi(x)\to h(x)O_\xi(x)h^\dagger(x)$ with $h(x)$ defined previously describes an invariant hidden local $U(1)$ symmetry. Thus, the chiral symmetry $SU(2)_L\times U(1)_Y$ covariance of the unrotated fields has been transferred totally to the hidden symmetry $U(1)$ covariance of the rotated fields. We can further find combination of electroweak gauge fields $g_2\frac{\tau^a}{2}W^a_{\xi,\mu}(x)-g_1\frac{\tau^3}{2}B_{\xi,\mu}(x)\to h(x)[g_2\frac{\tau^a}{2}W^a_{\xi,\mu}(x)-g_1\frac{\tau^3}{2}B_{\xi,\mu}(x)]h^\dagger(x)$ transforms covariantly, while alternative combination $g_2\frac{\tau^a}{2}W^a_{\xi,\mu}(x)+g_1\frac{\tau^3}{2}B_{\xi,\mu}(x)\to h(x)[g_2\frac{\tau^a}{2}W^a_{\xi,\mu}(x)+g_1\frac{\tau^3}{2}B_{\xi,\mu}(x)-2i\partial_\mu]h^\dagger(x)$ transforms as the "gauge field" of the hidden local $U(1)$ symmetry.

With technique used in Ref.[11], the integration over technigulon fields in Eq.(15b) can be formally integrated out with help of full n-point Green's function of the G^{α}_{μ}-field $G^{\alpha_1\ldots\alpha_n}_{\mu_1\ldots\mu_n}$, thus Eq.(15b) after integration becomes

$$\begin{aligned} e^{iS_{\rm norm}[U,W^a_\mu,B_\mu]} &= \int\mathcal{D}\bar\psi_\xi\mathcal{D}\psi_\xi\,\mathcal{F}[O_\xi]\delta(O_\xi-O^\dagger_\xi)\exp\Bigg[i\int d^4x\bar\psi_\xi\bigg(i\partial\!\!\!/-g_2\frac{\tau^a}{2}\slashed{W}^a_\xi P_L-g_1\frac{\tau^3}{2}\slashed{B}_\xi P_R\bigg)\psi_\xi \\ &\quad+\sum_{n=2}^{\infty}\int d^4x_1\ldots d^4x_n\frac{(-ig_{\rm TC})^n}{n!}G^{\alpha_1\ldots\alpha_n}_{\mu_1\ldots\mu_n}(x_1,\ldots,x_n)J^{\mu_1}_{\xi,\alpha_1}(x_1)\ldots J^{\mu_n}_{\xi,\alpha_n}(x_n)\Bigg]\,. \end{aligned} \tag{16}$$

where effective sources $J^\mu_{\xi,\alpha}(x)$ are identified as $J^\mu_{\xi,\alpha}(x)\equiv\bar\psi_\xi(x)t^\alpha\gamma^\mu\psi_\xi(x)$.

1. *Schwinger-Dyson Equation for Techniquark Propagator*

To show that the technicolor interaction indeed induces the condensate $\langle\bar\psi\psi\rangle\neq0$ which triggers the electroweak symmetry breaking, we investigate the behavior of the techniquark propagator $S^{\sigma\rho}(x,x')\equiv\langle\psi^\sigma_\xi(x)\bar\psi^\rho_\xi(x')\rangle$ in the following. *Neglecting the factor* $\mathcal{F}[O_\xi]\delta(O_\xi-O^\dagger_\xi)$ *in Eq.*(16), the total functional derivative of the integrand with respect to $\bar\psi^\sigma_\xi(x)$ is zero, (here and henceforth the suffixes σ and ρ are short notations for Lorentz spinor, techniflavor and technicolor indices,) *i.e.*,

$$\begin{aligned} 0 &= \int\mathcal{D}\bar\psi_\xi\mathcal{D}\psi_\xi\frac{\delta}{\delta\bar\psi^\sigma_\xi(x)}\exp\Bigg[\int d^4x(\bar\psi_\xi I+\bar I\psi_\xi)+i\int d^4x\bar\psi_\xi\bigg(i\partial\!\!\!/-g_2\frac{\tau^a}{2}\slashed{W}^a_\xi P_L-g_1\frac{\tau^3}{2}\slashed{B}_\xi P_R\bigg)\psi_\xi \\ &\quad+\sum_{n=2}^{\infty}\int d^4x_1\ldots d^4x_n\frac{(-ig_{\rm TC})^n}{n!}G^{\alpha_1\ldots\alpha_n}_{\mu_1\ldots\mu_n}(x_1,\ldots,x_n)J^{\mu_1}_{\xi,\alpha_1}(x_1)\ldots J^{\mu_n}_{\xi,\alpha_n}(x_n)\Bigg]\,, \end{aligned} \tag{17}$$

where $I(x)$ and $\bar I(x)$ are the external sources for, respectively, $\bar\psi_\xi(x)$ and $\psi_\xi(x)$; and which leads to

$$\begin{aligned} 0 &= \Bigg\langle\!\!\Bigg\langle I_\sigma(x)+i\bigg[i\partial\!\!\!/_x-g_2\frac{\tau^a}{2}\slashed{W}^a_\xi(x)P_L-g_1\frac{\tau^3}{2}\slashed{B}_\xi(x)P_R\bigg]_{\sigma\tau}\psi^\tau_\xi(x) \\ &\quad+\sum_{n=2}^{\infty}\int d^4x_2\ldots d^4x_n\frac{(-ig_{\rm TC})^n}{(n-1)!}G^{\alpha_1\ldots\alpha_n}_{\mu_1\ldots\mu_n}(x,x_2,\ldots,x_n)(t^{\alpha_1}\gamma^{\mu_1})_{\sigma\tau}\psi^\tau_\xi(x)J^{\mu_2}_{\xi,\alpha_2}(x_2)\ldots J^{\mu_n}_{\xi,\alpha_n}(x_n)\Bigg\rangle\!\!\Bigg\rangle_I\,, \end{aligned} \tag{18}$$

where we have defined the notation $\langle\langle\cdots\rangle\rangle_I$ in this section by

$$\begin{aligned} \langle\langle\mathcal{O}(x)\rangle\rangle_I &\equiv \int\mathcal{D}\bar\psi_\xi\mathcal{D}\psi_\xi\,\mathcal{O}(x)\exp\Bigg[\int d^4x(\bar\psi_\xi I+\bar I\psi_\xi) \\ &\quad+i\int d^4x\bar\psi_\xi\bigg(i\partial\!\!\!/-g_2\frac{\tau^a}{2}\slashed{W}^a_\xi P_L-g_1\frac{\tau^3}{2}\slashed{B}_\xi P_R\bigg)\psi_\xi \\ &\quad+\sum_{n=2}^{\infty}\int d^4x_1\ldots d^4x_n\frac{(-ig_{\rm TC})^n}{n!}G^{\alpha_1\ldots\alpha_n}_{\mu_1\ldots\mu_n}(x_1,\ldots,x_n)J^{\mu_1}_{\xi,\alpha_1}(x_1)\ldots J^{\mu_n}_{\xi,\alpha_n}(x_n)\Bigg] \end{aligned} \tag{19}$$

Taking functional derivative of Eq.(18) with respect to $I_\rho(y)$, and subsequently setting $I = \bar{I} = 0$, we obtain

$$\begin{aligned}
0 \;=\; & \delta_{\sigma\rho}\delta(x-y) + i\Big[i\not{\partial}_x - g_2\frac{\tau^a}{2}\not{W}^a_\xi(x)P_L - g_1\frac{\tau^3}{2}\not{B}_\xi(x)P_R\Big]_{\sigma\tau}\langle\psi^\tau_\xi(x)\bar{\psi}^\rho_\xi(y)\rangle \\
& -\sum_{n=2}^{\infty}\int d^4x_2\ldots d^4x_n\frac{(-ig_{\rm TC})^n}{(n-1)!}G^{\alpha_1\ldots\alpha_n}_{\mu_1\ldots\mu_n}(x,x_2,\ldots,x_n) \\
& \times\big\langle\bar{\psi}^\rho_\xi(y)(t^{\alpha_1}\gamma^{\mu_1})_{\sigma\tau}\psi^\tau_\xi(x)J^{\mu_2}_{\xi,\alpha_2}(x_2)\ldots J^{\mu_n}_{\xi,\alpha_n}(x_n)\big\rangle\;, \qquad (20)
\end{aligned}$$

where we have defined vacuum expectation value (VEV) $\langle\cdots\rangle$ by $\big\langle\mathcal{O}(x)\big\rangle \equiv \big\langle\!\big\langle\mathcal{O}(x)\big\rangle\!\big\rangle_I/\langle\langle\,1\,\rangle\rangle_I|_{I=\bar{I}=0}$. If we *neglect higher-point Green's functions*,and further *taking factorization approximation*, *i.e.*, $\big\langle\bar{\psi}^\rho_\xi(y)\psi^\tau_\xi(x)\bar{\psi}^\gamma_\xi(x_2)\psi^\delta_\xi(x_2)\big\rangle \approx \big\langle\bar{\psi}^\rho_\xi(y)\psi^\tau_\xi(x)\big\rangle\big\langle\bar{\psi}^\gamma_\xi(x_2)\psi^\delta_\xi(x_2)\big\rangle - \big\langle\bar{\psi}^\rho_\xi(y)\psi^\delta_\xi(x_2)\big\rangle\big\langle\bar{\psi}^\gamma_\xi(x_2)\psi^\tau_\xi(x)\big\rangle$, we obtain

$$\begin{aligned}
0 \;=\; & \delta_{\sigma\rho}\delta(x-y) + i\Big[i\not{\partial}_x - g_2\frac{\tau^a}{2}\not{W}^a_\xi(x)P_L - g_1\frac{\tau^3}{2}\not{B}_\xi(x)P_R\Big]_{\sigma\tau}\langle\psi^\tau_\xi(x)\bar{\psi}^\rho_\xi(y)\rangle \\
& -g^2_{\rm TC}\int d^4x_2\,G^{\alpha_1\alpha_2}_{\mu_1\mu_2}(x,x_2)(t^{\alpha_1}\gamma^{\mu_1})_{\sigma\tau}(t^{\alpha_2}\gamma^{\mu_2})_{\gamma\delta}\big\langle\bar{\psi}^\rho_\xi(y)\psi^\delta_\xi(x_2)\big\rangle\big\langle\bar{\psi}^\gamma_\xi(x_2)\psi^\tau_\xi(x)\big\rangle\;, \quad (21)
\end{aligned}$$

where we have used $\big\langle\bar{\psi}_\xi(x_2)t^{\alpha_2}\gamma^{\mu_2}\psi_\xi(x_2)\big\rangle = 0$, which comes from the Lorentz and gauge invariance of vacuum. We denote the technifermion propagator $S^{\sigma\rho}(x,x') \equiv \langle\psi^\sigma_\xi(x)\bar{\psi}^\rho_\xi(x')\rangle$, multiplying inverse of technifermion propagator in both sides of Eq.(21), it then is written as the Schwinger-Dyson equation (SDE) for techniquark propagator,

$$\begin{aligned}
0 \;=\; & S^{-1}_{\sigma\rho}(x,y) + i\Big[i\not{\partial}_x - g_2\frac{\tau^a}{2}\not{W}^a_\xi(x)P_L - g_1\frac{\tau^3}{2}\not{B}_\xi(x)P_R\Big]_{\sigma\rho}\delta(x-y) \\
& -g^2_{\rm TC}G^{\alpha_1\alpha_2}_{\mu_1\mu_2}(x,y)\Big[t^{\alpha_1}\gamma^{\mu_1}S(x,y)t^{\alpha_2}\gamma^{\mu_2}\Big]_{\sigma\rho}\;. \qquad (22)
\end{aligned}$$

By defining techniquark self energy $i\Sigma$ as

$$i\Sigma_{\sigma\rho}(x,y) \equiv S^{-1}_{\sigma\rho}(x,y) + i\Big[i\not{\partial}_x - g_2\frac{\tau^a}{2}\not{W}^a_\xi(x)P_L - g_1\frac{\tau^3}{2}\not{B}_\xi(x)P_R\Big]_{\sigma\rho}\delta(x-y)\;, \qquad (23)$$

the SDE (22) can also be written as

$$i\Sigma_{\sigma\rho}(x,y) = g^2_{\rm TC}G^{\alpha_1\alpha_2}_{\mu_1\mu_2}(x,y)\Big[t^{\alpha_1}\gamma^{\mu_1}S(x,y)t^{\alpha_2}\gamma^{\mu_2}\Big]_{\sigma\rho}\;. \qquad (24)$$

Moreover, from the fact that technigluon propagator is diagonal in the adjoint representation space of technicolor group, *i.e.*, $G^{\alpha\beta}_{\mu\nu}(x,y) = \delta^{\alpha\beta}G_{\mu\nu}(x,y)$, and techniquark propagator is diagonal in the techniquark representation space of technicolor group, and also $(t^\alpha t^\alpha)_{ab} =$

$C_2(N)\delta_{ab}$ for the fundamental representation of $SU(N)$, Eq.(24) is diagonal in technicolor indices a, b and diagonal part satisfy

$$i\Sigma^{ij}_{\eta\zeta}(x,y) = C_2(N)g^2_{\rm TC}G_{\mu_1\mu_2}(x,y)[\gamma^{\mu_1}S(x,y)\gamma^{\mu_2}]^{ij}_{\eta\zeta}\ , \tag{25}$$

where $\{i,\ j\}$, and $\{\eta,\ \zeta\}$ are, respectively, techniflavor and Lorentz spinor indices; and the Casimir operator $C_2(N) = (N^2-1)/(2N)$.

♣ $B_{\xi,\mu} = W^a_{\xi,\mu} = 0$ Case: the Gap Equation

The technigluon propagator in Landau gauge is $G^{\alpha\beta}_{\mu\nu}(x,y) = \delta^{\alpha\beta}\int\frac{d^4p}{(2\pi)^4}e^{-ip(x-y)}G_{\mu\nu}(p^2)$ with $G_{\mu\nu}(p^2) = \frac{i}{-p^2[1+\Pi(-p^2)]}(g_{\mu\nu}-p_\mu p_\nu/p^2)$. And the techniquark self energy and propagator are respectively

$$\Sigma^{ij}_{\eta\zeta}(x,y) = \int\frac{d^4p}{(2\pi)^4}e^{-ip(x-y)}\,\Sigma^{ij}_{\eta\zeta}(-p^2) \qquad S^{ij}_{\eta\zeta}(x,y) = \int\frac{d^4p}{(2\pi)^4}e^{-ip(x-y)}S^{ij}_{\eta\zeta}(p)\ , \tag{26}$$

with $S^{ij}_{\eta\zeta}(p) = i[1/(\not p - \Sigma(-p^2)]^{ij}_{\eta\zeta}$. Substituting above results into the SDE (25), we have

$$\Sigma^{ij}_{\eta\zeta}(-p^2) = \int\frac{d^4q}{(2\pi)^4}\frac{-C_2(N)g^2_{\rm TC}}{(p-q)^2[1+\Pi(-(p-q)^2)]}\left[g_{\mu\nu}-\frac{(p-q)_\mu(p-q)_\nu}{(p-q)^2}\right]\left[\gamma^\mu\frac{i}{\not q-\Sigma(-q^2)}\gamma^\nu\right]^{ij}_{\eta\zeta} \tag{27}$$

from which we can see that the solution of the techniquark self energy must be diagonal in techniflavor space, since the integration kernel is independent of techniflavor indices, *i.e.*, $\Sigma^{ij}_{\eta\zeta}(-p^2) = \delta^{ij}\Sigma_{\eta\zeta}(-p^2)$. With the assumption that the techniquark self energy is diagonal and equal in Lorentz spinor space, leads to the following two equations

$$i\,\Sigma(-p^2) = 3C_2(N)\int\frac{d^4q}{4\pi^3}\frac{\alpha_{\rm TC}[-(p-q)^2]}{(p-q)^2}\frac{\Sigma(-q^2)}{q^2-\Sigma^2(-q^2)}\ , \tag{28a}$$

$$0 = \int d^4q\,\frac{\alpha_{\rm TC}[-(p-q)^2]}{(p-q)^2}\left[g_{\mu\nu}-\frac{(p-q)_\mu(p-q)_\nu}{(p-q)^2}\right]\gamma^\mu\frac{\not q}{q^2-\Sigma^2(-q^2)}\gamma^\nu\ . \tag{28b}$$

In which we have labeled the integration kernel with running coupling constant $\alpha_{\rm TC}(-p^2) \equiv g^2_{\rm TC}(-p^2)/(4\pi) = g^2_{\rm TC}/(4\pi[1+\Pi(-p^2)])$. Eq.(28b) is automatically satisfied when taken *the approximation*, $\alpha_{\rm TC}[(p_E-q_E)^2] = \alpha_{\rm TC}(p_E^2)\theta(p_E^2-q_E^2)+\alpha_{\rm TC}(q_E^2)\theta(q_E^2-p_E^2)$. The other equation (28a) can be written in Euclidean space as

$$\Sigma(p_E^2) = 3C_2(N)\int\frac{d^4q_E}{4\pi^3}\frac{\alpha_{\rm TC}[(p_E-q_E)^2]}{(p_E-q_E)^2}\frac{\Sigma_(q_E^2)}{q_E^2+\Sigma^2(q_E^2)}\ , \tag{29}$$

If there is nonzero solution for above equation, we will obtain nonzero techniquark condensate $\langle\bar\psi^k_\xi\psi^j_\xi\rangle$ with k and j techniflavor indices,

$$\langle\bar\psi^k_\xi(x)\psi^j_\xi(x)\rangle\ =\ -{\rm tr}_{lc}[S^{jk}(x,x)] = -4N\delta^{jk}\int\frac{d^4p_E}{(2\pi)^4}\frac{\Sigma(p_E^2)}{p_E^2+\Sigma^2(p_E^2)}\ , \tag{30}$$

where tr_{lc} is the trace with respect to Lorentz, technicolor indices, and where we have used that the techniquark self energy, the solution of the SDE, must be diagonal in techniflavor space. Thus, nonzero techniquark self energy can give a nontrivial diagonal condensate $\langle\bar{\psi}\psi\rangle \neq 0$, which spontaneously breaks $SU(2)_L \times U(1)_Y$ to $U(1)_{\mathrm{em}}$.

♣ $B_{\xi,\mu} \neq 0$ and $W^a_{\xi,\mu} \neq 0$ Case: the Lowest-order Approximation

In the following we consider the effects of the nonzero electroweak gauge fields $B_{\xi,\mu}$ and $W^a_{\xi,\mu}$. The SDE (25) is explicitly

$$\begin{aligned}\Sigma(x,y) &= C_2(N)g^2_{\mathrm{TC}}G_{\mu\nu}(x,y)\\ &\times\gamma^\mu\left[\left(i\partial\!\!\!/_x - g_2\frac{\tau^a}{2}W\!\!\!\!/^a_\xi(x)P_L - g_1\frac{\tau^3}{2}B\!\!\!/_\xi(x)P_R\right)\delta(x-y) - \Sigma(x,y)\right]^{-1}\gamma^\nu\ , \quad (31)\end{aligned}$$

where the techniflavor and Lorentz spinor indices of the techniquark self energy are implicitly contained. In this case, the self energy can no longer be written as the function on the derivatives with respect to spacetime, *i.e.*, $\Sigma(x,y) \neq \Sigma(\partial^2_x)\delta(x-y)$.

Suppose the function $\Sigma(-p^2)$ is a solution of the SDE in the case $B_{\xi,\mu} = W^a_{\xi,\mu} = 0$, that is, it satisfies the equation

$$\Sigma(-p^2) = C_2(N)g^2_{\mathrm{TC}}\int\frac{d^4q}{(2\pi)^4}G_{\mu\nu}(q^2)\gamma^\mu\frac{1}{q\!\!\!/ + p\!\!\!/ - \Sigma[-(q+p)^2]}\gamma^\nu\ , \quad (32)$$

where in the second equality the legality of the integration measure translation comes from the logarithmical divergence of the fermion self energy. Replacing the variable p by $p+\Delta$ in Eq.(32) and subsequently integrating over p with the weight $e^{-ip(x-y)}$, we obtain, as long as Δ is commutative with ∂_x and Dirac matrices, Eq.(32) imply

$$\Sigma[(\partial_x - i\Delta)^2]\delta(x-y) = C_2(N)g^2_{\mathrm{TC}}G_{\mu\nu}(x,y)\gamma^\mu\frac{1}{i\partial\!\!\!/_x + \Delta\!\!\!/ - \Sigma[-(i\partial_x+\Delta)^2]}\delta(x-y)\gamma^\nu\ . \quad (33)$$

Even if Δ is noncommutative with ∂_x and Dirac matrices, the above equation holds as the lowest order approximation, for the commutator $[\partial\!\!\!/, \Delta]$ is higher order of momentum than Δ itself. Now if we take Δ to be $-g_2\frac{\tau^a}{2}W^a_\xi P_L - g_1\frac{\tau^3}{2}B_\xi P_R$, *ignoring its noncommutative property with ∂_x and Dirac matrices*, Eq.(33) is just the SDE (31) in the case $B_{\xi,\mu} \neq 0$ and $W^a_{\xi,\mu} \neq 0$. Thus, $\Sigma[(\partial^x_\mu + ig_2\frac{\tau^a}{2}W^a_{\xi,\mu}P_L + ig_1\frac{\tau^3}{2}B_{\xi,\mu}P_R)^2]\delta(x-y)$, which is the hidden symmetry $U(1)$ covariant, can be regarded as the lowest-order solution of Eq.(31). To further simplify the calculations and still keep this hidden-symmetry covariance of the self energy, we can *reduce the covariant derivative inside the self energy* $\nabla_\mu \equiv \partial_\mu + ig_2\frac{\tau^a}{2}W^a_{\xi,\mu}P_L + ig_1\frac{\tau^3}{2}B_{\xi,\mu}P_R$ *to its minimal-coupling form*

$$\overline{\nabla}_\mu \equiv \partial_\mu + \frac{i}{2}[g_2\frac{\tau^a}{2}W^a_{\xi,\mu}(x) + g_1\frac{\tau^3}{2}B_{\xi,\mu}(x)] \; . \tag{34}$$

In which as we mentioned before $[g_2\frac{\tau^a}{2}W^a_{\xi,\mu}(x) + g_1\frac{\tau^3}{2}B_{\xi,\mu}(x)]/2$ transforms as a gauge field under hidden $U(1)$ symmetry transformations. Thus, if the function $\Sigma(\partial_x^2)\delta(x-y)$ is the self-energy solution of the SDE in the case $B_{\xi,\mu} = W^a_{\xi,\mu} = 0$, we can replace its argument ∂_x by the minimal-coupling covariant derivative $\overline{\nabla}_x$, *i.e.*, $\Sigma(\overline{\nabla}_x^2)\delta(x-y)$, as an approximate solution of the SDE in the case $B_{\xi,\mu} \neq 0$ and $W^a_{\xi,\mu} \neq 0$.

2. *Effective Action*

The exponential terms on the right-hand side of Eq.(16) can be written explicitly as

$$\begin{aligned}&\sum_{n=2}^{\infty}\int d^4x_1\ldots d^4x_n\frac{(-ig_{\rm TC})^n}{n!}G^{\alpha_1\ldots\alpha_n}_{\mu_1\ldots\mu_n}(x_1,\ldots,x_n)J^{\mu_1}_{\xi,\alpha_1}(x_1)\ldots J^{\mu_n}_{\xi,\alpha_n}(x_n)\\ &\approx \int d^4x d^4x'\bar{\psi}^\sigma_\xi(x)\Pi_{\sigma\rho}(x,x')\psi^\rho_\xi(x') \; ,\end{aligned} \tag{35}$$

where we have taken the approximation of *replacing the summation over* $2n$*-fermion interactions with parts of them by their vacuum expectation values*, that is,

$$\Pi_{\sigma\rho}(x,x') \;=\; \sum_{n=2}^{\infty}\Pi^{(n)}_{\sigma\rho}(x,x') \; , \tag{36}$$

$$\begin{aligned}\Pi^{(n)}_{\sigma\rho}(x,x') \;=\;& n\times\int d^4x_2\ldots d^4x_{n-1}\frac{(-ig_{\rm TC})^n}{n!}G^{\alpha_1\ldots\alpha_n}_{\mu_1\ldots\mu_n}(x,x_2\ldots,x_{n-1},x')\Big\langle(t^{\alpha_1}\gamma^{\mu_1})_{\sigma\sigma_1}\psi^{\sigma_1}_\xi(x)\\ &\times\bar{\psi}_\xi(x_2)t^{\alpha_2}\gamma^{\mu_2}\psi_\xi(x_2)\ldots\bar{\psi}_\xi(x_{n-1})t^{\alpha_{n-1}}\gamma^{\mu_{n-1}}\psi_\xi(x_{n-1})\bar{\psi}^{\rho_n}_\xi(x')(t^{\alpha_n}\gamma^{\mu_n})_{\rho_n\rho}\Big\rangle\end{aligned} \tag{37}$$

where the factor n comes from n different choices of unaveraged $\bar{\psi}_\xi\psi_\xi$, and the lowest term of which is

$$\Pi^{(2)}_{\sigma\rho}(x,x') \;=\; -g^2_{\rm TC}G^{\alpha_1\alpha_2}_{\mu_1\mu_2}(x,x')\Big[t^{\alpha_1}\gamma^{\mu_1}S(x,y)t^{\alpha_2}\gamma^{\mu_2}\Big]_{\sigma\rho} \; . \tag{38}$$

Comparing Eq.(38) with Eq.(24), we have

$$i\Pi^{(2)}_{\sigma\rho}(x,x') = \Sigma_{\sigma\rho}(x,x') \approx \Sigma_{\sigma\rho}(\overline{\nabla}^2_x)\delta(x-y) \; . \tag{39}$$

By neglecting the factor $\mathcal{F}[O_\xi]\delta(O_\xi - O_\xi^\dagger)$ *in Eq.*(16), we have

$$\begin{aligned}
e^{iS_{\rm norm}[U,W^a_\mu,B_\mu]} &\approx \int \mathcal{D}\bar{\psi}_\xi\mathcal{D}\psi_\xi \exp\Big[i\int d^4x\bar{\psi}_\xi\Big(i\not\partial - g_2\frac{\tau^a}{2}\not W^a_\xi P_L - g_1\frac{\tau^3}{2}\not B_\xi P_R\Big)\psi_\xi \\
&\qquad + \int d^4x d^4x'\bar{\psi}^\sigma_\xi(x)\Pi_{\sigma\rho}(x,x')\psi^\rho_\xi(x')\Big] \\
&\approx \mathrm{Det}[i\not\partial - g_2\frac{\tau^a}{2}\not W^a_\xi P_L - g_1\frac{\tau^3}{2}\not B_\xi P_R - \Sigma(\overline{\nabla}^2)] \qquad (40)
\end{aligned}$$

where in the second equality we have taken further approximation of *keeping only the lowest order, i.e.* $\Pi^{(2)}_{\sigma\rho}(x,x')$, *of* $\Pi_{\sigma\rho}(x,x')$. With these three approximations, we have

$$S_{\rm norm}[U,W^a_\mu,B_\mu] = -i\mathrm{Tr}\log[i\not\partial - g_2\frac{\tau^a}{2}\not W^a_\xi P_L - g_1\frac{\tau^3}{2}\not B_\xi P_R - \Sigma(\overline{\nabla}^2)]\;, \qquad (41)$$

As done in [12], we can parameterize the normal part of the effective action as follows

$$\begin{aligned}
&S_{\rm norm}[U,W^a_\mu,B_\mu] = -i\mathrm{Tr}\log[i\not\partial + \not v + \not a\gamma_5 - \Sigma(\overline{\nabla}^2)] \\
&= \int d^4x\,\mathrm{tr}_f\Big[(F_0^{\rm 1D})^2a^2 - \mathcal{K}_1^{\rm 1D}(d_\mu a^\mu)^2 - \mathcal{K}_2^{\rm 1D}(d_\mu a_\nu - d_\nu a_\mu)^2 + \mathcal{K}_3^{\rm 1D}(a^2)^2 + \mathcal{K}_4^{\rm 1D}(a_\mu a_\nu)^2 \\
&\quad -\mathcal{K}_{13}^{\rm 1D}V_{\mu\nu}V^{\mu\nu} + i\mathcal{K}_{14}^{\rm 1D}a_\mu a_\nu V^{\mu\nu}\Big] + \mathcal{O}(p^6)\;, \qquad (42)
\end{aligned}$$

where the fields v_μ, a_μ are identified with $v_\mu \equiv -\frac{1}{2}(g_2\frac{\tau^a}{2}W^a_{\xi,\mu} + g_1\frac{\tau^3}{2}B_{\xi,\mu})$ and $a_\mu \equiv \frac{1}{2}(g_2\frac{\tau^a}{2}W^a_{\xi,\mu} - g_1\frac{\tau^3}{2}B_{\xi,\mu})$ and $d_\mu a_\nu \equiv \partial_\mu a_\nu - i[v_\mu, a_\nu]$, $V_{\mu\nu} \equiv i[\partial_\mu - iv_\mu, \partial_\nu - iv_\nu]$. $\mathcal{K}_i^{\rm 1D}$ coefficients with superscript 1D to denote present one doublet model are functions of techniquark self energy $\Sigma(p^2)$ and detail expressions are already written down in (36) of Ref.[12] with the replacement of $N_c \to N$.

For anomaly part, compare (15c) and (41), we find its U field dependent part can be produced from normal part by vanishing techniquark self energy Σ, i.e.

$$iS_{\rm anom}[U,W^a_\mu,B_\mu] = \mathrm{Tr}\log(i\not\partial - g_2\frac{\tau^a}{2}\not W^a P_L - g_1\frac{\tau^3}{2}\not B P_R) - iS_{\rm norm}[U,W^a_\mu,B_\mu]|_{\Sigma=0} \qquad (43)$$

Notice that U field independent part of pure gauge field part is irrelevant to EWCL. Combine with (42), above relation imply

$$\begin{aligned}
iS_{\rm anom}[U,W^a_\mu,B_\mu] &= \mathrm{Tr}\log(i\not\partial - g_2\frac{\tau^a}{2}\not W^a P_L - g_1\frac{\tau^3}{2}\not B P_R) + i\int d^4x\,\mathrm{tr}_f\Big[-\mathcal{K}_1^{\rm 1D,(anom)}(d_\mu a^\mu)^2 \\
&\quad -\mathcal{K}_2^{\rm 1D,(anom)}(d_\mu a_\nu - d_\nu a_\mu)^2 + \mathcal{K}_3^{\rm 1D,(anom)}(a^2)^2 + \mathcal{K}_4^{\rm 1D,(anom)}(a_\mu a_\nu)^2 \\
&\quad -\mathcal{K}_{13}^{\rm 1D,(anom)}V_{\mu\nu}V^{\mu\nu} + i\mathcal{K}_{14}^{\rm 1D,(anom)}a_\mu a_\nu V^{\mu\nu}\Big] + \mathcal{O}(p^6)\;, \qquad (44)
\end{aligned}$$

with

$$\mathcal{K}_i^{\rm 1D,(anom)} = -\mathcal{K}_i^{\rm 1D}|_{\Sigma=0} \qquad\qquad i=1,2,3,4,13,14 \tag{45}$$

where we have used result that $F_0^{\rm 1D}|_{\Sigma=0}=0$. Combine normal and anomaly part contribution together, with help of (15a), we finally find

$$\begin{aligned}
S_{\rm eff}[U,W_\mu^a,B_\mu] \;=\; & -\frac{1}{4}\int d^4x(W_{\mu\nu}^a W^{a,\mu\nu}+B_{\mu\nu}B^{\mu\nu})+{\rm Tr}\log(i\partial\!\!\!/-g_2\frac{\tau^a}{2}W\!\!\!\!/^{\,a}P_L-g_1\frac{\tau^3}{2}B\!\!\!/P_R)\\
& +i\int d^4x\,{\rm tr}_f\bigg[(F_0^{\rm 1D})^2a^2-\mathcal{K}_1^{\rm 1D,\Sigma\neq0}(d_\mu a^\mu)^2-\mathcal{K}_2^{\rm 1D,\Sigma\neq0}(d_\mu a_\nu-d_\nu a_\mu)^2+\mathcal{K}_3^{\rm 1D,\Sigma\neq0}(a^2)^2\\
& +\mathcal{K}_4^{\rm 1D,\Sigma\neq0}(a_\mu a_\nu)^2-\mathcal{K}_{13}^{\rm 1D,\Sigma\neq0}V_{\mu\nu}V^{\mu\nu}+i\mathcal{K}_{14}^{\rm 1D,\Sigma\neq0}a_\mu a_\nu V^{\mu\nu}\bigg]+\mathcal{O}(p^6)\;,
\end{aligned}\tag{46}$$

with $\mathcal{K}_i^{\rm 1D,\Sigma\neq0}$ be Σ dependent part of $\mathcal{K}_i$

$$\mathcal{K}_i^{\rm 1D,\Sigma\neq0}\equiv\mathcal{K}_i^{\rm 1D}-\mathcal{K}_i^{\rm 1D}|_{\Sigma=0} \qquad\qquad i=1,2,3,4,13,14 \tag{47}$$

After some algebras, terms in Eq.(46) can be reexpressed in terms of X_μ and $\overline{W}_{\mu\nu}$ which are just standard EWCL given in Ref.[2] with coefficients

$$\begin{aligned}
& f^2=(F_0^{\rm 1D})^2\;,\qquad \beta_1=0\;,\qquad \alpha_1=\frac{\mathcal{K}_2^{\rm 1D,\Sigma\neq0}-\mathcal{K}_{13}^{\rm 1D,\Sigma\neq0}}{2}\;,\qquad \alpha_2=\alpha_3=-\frac{\mathcal{K}_{13}^{\rm 1D,\Sigma\neq0}}{4}+\frac{\mathcal{K}_{14}^{\rm 1D,\Sigma\neq0}}{16}\;,\\
& \alpha_4=\frac{\mathcal{K}_4^{\rm 1D,\Sigma\neq0}+2\mathcal{K}_{13}^{\rm 1D,\Sigma\neq0}-\mathcal{K}_{14}^{\rm 1D,\Sigma\neq0}}{16}\;,\qquad \alpha_5=\frac{\mathcal{K}_3^{\rm 1D,\Sigma\neq0}-\mathcal{K}_4^{\rm 1D,\Sigma\neq0}-4\mathcal{K}_{13}^{\rm 1D,\Sigma\neq0}+2\mathcal{K}_{14}^{\Sigma\neq0}}{32}\;.
\end{aligned}\tag{48}$$

With formulae of $\mathcal{K}_i$ coefficients given in Ref.[12], we can substitute the solution of SD equation (29) into them and then obtain numerical results for those nonzero α_i coefficients. In TABLE I, we list down the numerical calculation results for different kind of dynamics.

To obtain above numerical result, we have solved Schwinger-Dyson equation (29) with following running coupling which was used as model A in Ref.[12]

$$\alpha_{\rm TC}(p^2)=\begin{cases}7\frac{12\pi}{(11N-2N_f)}, & \text{for }\ln(p^2/\Lambda_{\rm TC}^2)\leq-2;\\ \{7-\frac{4}{5}[2+\ln(p^2/\Lambda_{\rm TC}^2)]^2\}\frac{12\pi}{(11N-2N_f)}, & \text{for }-2\leq\ln(p^2/\Lambda_{\rm TC}^2)\leq0.5;\\ \dfrac{1}{\ln(p^2/\Lambda_{\rm TC}^2)}\dfrac{12\pi}{(11N-2N_f)}, & \text{for }0.5\leq\ln(p^2/\Lambda_{\rm TC}^2)\end{cases}\tag{49}$$

In which the fermion number is taken to be $N_f=2$ corresponding to present one doublet techniquark. Although there is a dimensional parameter $\Lambda_{\rm TC}$ appear in $\alpha_{\rm TC}(p^2)$, except dimensional coefficient $F_0^{\rm 1D}$, all dimensionless result coefficients $\alpha_i,\;\; i=1,2,3,4,5$ are independent of this parameter. This can be seen as follows, if we scale up $\Lambda_{\rm TC}$ as $\lambda\Lambda_{\rm TC}$,

TABLE I. The obtained nonzero values of the $O(p^4)$ coefficients $\alpha_1, \alpha_2 = \alpha_3, \alpha_4, \alpha_5$ for one doublet technicolor model with the conventional strong interaction QCD theory values given in Ref.[12] for model A and experimental values for comparison. $\Lambda_{\rm TC}$ and $\Lambda_{\rm QCD}$ are in TeV, they are determined by $f = 250$GeV and $f_\pi = 93$MeV respectively. The coefficients are in units of 10^{-3}. QCD values are taken by using relation (9).

N	$\Lambda_{\rm TC}$	α_1	$\alpha_2 = \alpha_3$	α_4	α_5
3	1.34	-6.90	-2.43	2.02	-2.69
4	1.15	-9.26	-3.28	2.87	-3.69
5	1.03	-11.6	-4.11	3.60	-4.62
6	0.94	-13.9	-4.93	4.32	-5.54
QCD Theor	$\Lambda_{\rm QCD}$=0.484*10^{-3}	-7.06	-2.54	2.20	-2.81
QCD Expt		-6.0 ± 0.7	-2.7 ± 0.4	1.7 ± 0.7	-1.3 ± 1.5

$\alpha_{\rm TC}(p)$ defined above satisfy $\alpha_{\rm TC}(p^2)|_{\lambda\Lambda_{\rm TC}} = \alpha_{\rm TC}(\lambda^{-2}p^2)|_{\Lambda_{\rm TC}}$ which, by (29), result a scaling-up techniquark self energy $\Sigma(p^2)|_{\lambda\Lambda_{\rm TC}} = \lambda\Sigma(\lambda^{-2}p^2)|_{\Lambda_{\rm TC}}$, since an alternative expression of (29) is

$$\lambda\Sigma(\lambda^{-2}p_E^2) = 3C_2(N)\int\frac{d^4q_E}{4\pi^3}\frac{\alpha_{\rm TC}[\lambda^{-2}(p_E-q_E)^2]}{(p_E-q_E)^2}\frac{\lambda\Sigma(\lambda^{-2}q_E^2)}{q_E^2+\lambda^2\Sigma^2(\lambda^{-2}q_E^2)}\ . \tag{50}$$

Further from (36) of Ref.[12], we find coefficients $\mathcal{K}_i^{\Sigma\neq0},\ \ i = 1,2,3,4,13,14$ are invariant and F_0 is changed to λF_0 under exchanging $\Sigma(p^2) \to \lambda\Sigma(\lambda^{-2}p^2)$ if we take cutoff in the formulae $\Lambda \to \infty$. Due to this invariance for $\mathcal{K}_i^{\Sigma\neq0},\ \ i = 1,2,3,4,13,14$, from (51), we can see then $\alpha_i,\ \ i = 1,2,3,4,5$ are independent of $\Lambda_{\rm TC}$ and $F_0^{\rm 1D}$ scales same as $\Lambda_{\rm TC}$. It is this scale dependence for F_0 which makes ordinary QCD contribution small to electroweak symmetry breaking and leads necessity for new interactions at higher energy scale. The scale relations above are result of present rough approximations, it will simplify our future computations very much in next section.

From TABLE I, we see that within error of our approximation, the numerical result exhibit the scaling-up behavior among different N.

C. Comparison and Discussion on Two Prescriptions

Compare results from Gasser-Leutwyler's prescription and dynamical computation prescription, (9) and (48), we find results are same as long as we identify

$$\begin{aligned}
&H_1^{\mathrm{1D}}=-\frac{\mathcal{K}_2^{\mathrm{1D},\Sigma\neq0}+\mathcal{K}_{13}^{\mathrm{1D},\Sigma\neq0}}{4} \qquad L_{10}^{\mathrm{1D}}=\frac{\mathcal{K}_2^{\mathrm{1D},\Sigma\neq0}-\mathcal{K}_{13}^{\mathrm{1D},\Sigma\neq0}}{2}\\
&L_9^{\mathrm{1D}}=\frac{\mathcal{K}_{13}^{\mathrm{1D},\Sigma\neq0}}{2}-\frac{\mathcal{K}_{14}^{\mathrm{1D},\Sigma\neq0}}{8} \qquad L_2^{\mathrm{1D}}=\frac{\mathcal{K}_4^{\mathrm{1D},\Sigma\neq0}+2\mathcal{K}_{13}^{\mathrm{1D},\Sigma\neq0}-\mathcal{K}_{14}^{\mathrm{1D},\Sigma\neq0}}{16}\\
&L_1^{\mathrm{1D}}+\frac{L_3^{\mathrm{1D}}}{2}=\frac{\mathcal{K}_3^{\mathrm{1D},\Sigma\neq0}-\mathcal{K}_4^{\mathrm{1D},\Sigma\neq0}-4\mathcal{K}_{13}^{\mathrm{1D},\Sigma\neq0}+2\mathcal{K}_{14}^{\mathrm{1D},\Sigma\neq0}}{32}
\end{aligned} \tag{51}$$

which is just the result (25) obtained in Ref.[12]. This shows that two prescriptions are equivalent in results. The merit of Gasser-Leutwyler's prescription is its simplicity and express result coefficients of electroweak chiral lagrangian in terms those in Gasser-Leutwyler chiral lagrangian for pseudo scalar mesons in strong interaction, but we can only apply this prescription to so called QCD-like theories for which the technicolor interaction must be vector-like. On the other hand, dynamical computation prescription, although much more complex but touch the dynamics details, do not limit us in the type of detail interactions. this has very strong potential to be applied to more complicated theories, such as chiral-like technicolor models. Since we involve detail dynamical computation in this prescription, not like Gasser-Leutwyler's prescription the coefficients are expressed in terms of strong interaction experiment fixed values, we can give detail theoretical computation result for all coefficients and it further allow us to test possible effects on the coefficients from variation of the dynamics.

The first property qualitatively drew out from (41) and (15c) for their trace operation is that all coefficients are proportional to N. This is the well known scaling-up result for one doublet technicolor model, i.e., present coefficients can be got by (9) but identify the L_i with the corresponding Gasser-Leutwyler chiral lagrangian for pseudo scalar mesons by an scaling-up factor N/N_c with $N_c=3$ be number of color for strong interaction. In fact, it was this direct correspondence lead to the death of one doublet technicolor model, since negative experiment value for L_{10}^{1D} result large positive $S=-16\pi\alpha_1=-16\pi L_{10}^{\mathrm{1D}}$ parameter which contradict with present electroweak precision measurement data.

The second property quantitatively drew out from (41) and (15c) is that except the overall N factor in front of all coefficients, remaining part of coefficients depends on dynamics, so exactly speaking, they are not precisely equivalent to their strong interaction partners. For one doublet technicolor model, when $N \neq N_c$, not only we will have overall scaling-up factor N/N_c, but also we will have different techniquark self energy Σ due to the difference in running coupling constant (49) appeared in SDE (29). $N_f = 2$ also cause similar differences. But, since the estimations over the values in original strong interaction suffers large errors either in experiment or theories, this difference caused by dynamics hides in these uncertainties.

III. DERIVATION OF THE ELECTROWEAK CHIRAL LAGRANGIAN FROM A TOPCOLOR-ASSISTED TECHNICOLOR MODEL

There are several options in topcolor-assisted technicolor model building: (1) TC breaks both the EW interactions and the TopC interactions; (2) TC breaks EW, and something else breaks TopC; (3) TC breaks only TopC and something else drives EWSB (*e.g.*, a fourth generation condensate driven by TopC). For definiteness, we will focus on a skeletal model in category (1) in the following.

Consider a schematic TC2 model proposed by C. T. Hill [21]. The technicolor group is chosen to be $G_{\rm TC} = SU(3)_{\rm TC1} \times SU(3)_{\rm TC2}$. The gauge charge assignments of techniquarks in $G_{\rm TC} \times SU(3)_1 \times SU(3)_2 \times SU(2)_L \times U(1)_{Y_1} \times U(1)_{Y_2}$ are shown as Table II.

TABLE II. Gauge charge assignments of techniquarks a schematic Topcolor-assisted Technicolor model. Ordinary quarks and additional fields (such as leptons) required for anomaly cancellation are not shown. The techniquark condensate $\langle \bar{Q}Q \rangle$ breaks $SU(3)_1 \times SU(3)_2 \times U(1)_{Y_1} \times U(1)_{Y_2} \to SU(3) \times U(1)_Y$, while $\langle \bar{T}T \rangle$ breaks $SU(2)_L \times U(1)_Y \to U(1)_{\rm EM}$.

field	$SU(3)_{\rm TC1}$	$SU(3)_{\rm TC2}$	$SU(3)_1$	$SU(3)_2$	$SU(2)_L$	$U(1)_{Y_1}$	$U(1)_{Y_2}$
Q_L	3	1	3	1	1	$\frac{1}{2}$	0
Q_R	3	1	1	3	1	0	$\frac{1}{2}$
$T_L = (T,\ B)_L$	1	3	1	1	2	0	$\frac{1}{6}$
$T_R = (T,\ B)_R$	1	3	1	1	1	0	$(\frac{2}{3},\ -\frac{1}{3})$

The action of the symmetry breaking sector (SBS) then is

$$S_{\rm SBS} = \int d^4x(\mathcal{L}_{\rm gauge} + \mathcal{L}_{\rm techniquark}) \; , \tag{52}$$

with different part of Lagrangian given by

$$\begin{aligned}
\mathcal{L}_{\rm gauge} &= -\frac{1}{4}F^{\alpha}_{1\mu\nu}F^{\alpha\mu\nu}_{1} - \frac{1}{4}F^{\alpha}_{2\mu\nu}F^{\alpha\mu\nu}_{2} - \frac{1}{4}A^{A}_{1\mu\nu}A^{A\mu\nu}_{1} - \frac{1}{4}A^{A}_{2\mu\nu}A^{A\mu\nu}_{2} - \frac{1}{4}W^{a}_{\mu\nu}W^{a\mu\nu} \\
&\quad -\frac{1}{4}B_{1\mu\nu}B^{\mu\nu}_{1} - \frac{1}{4}B_{2\mu\nu}B^{\mu\nu}_{2} \; , \qquad (53) \\
\mathcal{L}_{\rm techniquark} &= \bar{Q}\big(i\partial\!\!\!/ - g_{31}r^{\alpha}_{1}G\!\!\!/^{\alpha}_{1} - h_1\frac{\lambda^A}{2}A\!\!\!/^{A}_{1}P_L - h_2\frac{\lambda^A}{2}A\!\!\!/^{A}_{2}P_R - q_1\frac{1}{2}B\!\!\!/_1P_L - q_2\frac{1}{2}B\!\!\!/_2P_R\big)Q \\
&\quad +\bar{T}\big[i\partial\!\!\!/ - g_{32}r^{\alpha}_{2}G\!\!\!/^{\alpha}_{2} - g_2\frac{\tau^a}{2}W\!\!\!\!/\,P_L - q_2\frac{1}{6}B\!\!\!/_2P_L - q_2(\frac{1}{6}+\frac{\tau^3}{2})B\!\!\!/_2P_R\big]T \; , \qquad (54)
\end{aligned}$$

where g_{31}, g_{32}, h_1, h_2, g_2, q_1 and q_2 are the coupling constants of, respectively, $SU(3)_{\rm TC1}$, $SU(3)_{\rm TC2}$, $SU(3)_1$, $SU(3)_2$, $SU(2)_L$, $U(1)_{Y_1}$ and $U(1)_{Y_2}$; and the corresponding gauge fields (field strength tensors) are denoted by $G^{\alpha}_{1\mu}$, $G^{\alpha}_{2\mu}$, $A^{A}_{1\mu}$, $A^{A}_{2\mu}$, W^{a}_{μ}, $B_{1\mu}$ and $B_{2\mu}$ ($F^{\alpha}_{1\mu\nu}$, $F^{\alpha}_{2\mu\nu}$, $A^{A}_{1\mu\nu}$, $A^{A}_{2\mu\nu}$, $W^{a}_{\mu\nu}$, $B_{1\mu\nu}$ and $B_{2\mu\nu}$) with the superscripts α and A running from 1 to 8 and a from 1 to 3; r^{α}_{1} and r^{α}_{2} ($\alpha = 1,\ldots,8$) are the generators of, respectively, $SU(3)_{\rm TC1}$ and $SU(3)_{\rm TC2}$, while λ^A ($A = 1,\ldots,8$) and τ^a ($a = 1,2,3$) are, respectively, Gell-Mann and Pauli matrices. We do not consider the ordinary quarks in this work for following considerations, as we mentioned in the Introduction that this paper only discuss bosonic part of EWCL and matter part of EWCL will discussed in future. The matter part of EWCL mainly deals with effective interactions among ordinary fermions which certainly include ordinary quarks. Ignoring discussion of these effective interactions, only concentrate on their contribution to bosonic part EWCL coefficients is not self-consistent and efficient. Further one special feature of topcolor-assisted technicolor model is its arrangements on the interactions among ordinary quarks, especially for top and bottom quark mass splitting and the top pions resulted from top quark condensation through topcolor interactions, therefore dealing with quark interactions is a separated important issue which needs special care, previous formal derivations from underlying gauge theory to low energy chiral Lagrangian, no matter QCD and electroweak theory, all not involve in matter part of chiral Lagrangian and further initial computation shows that we need some special techniques to handle top-bottom splitting which are beyond those techniques developed in this paper, to simplify the computations and reduce the lengthy formulae, we will not involve in discussion of ordinary quark in this paper and would rather specially focus our attentions on this issue in future works.

The strategy to derive the electroweak chiral Lagrangian from the schematic topcolor-assisted technicolor model can be formulated as

$$\begin{aligned}\exp\left(iS_{\mathrm{EW}}[W^a_\mu,B_\mu]\right) &= \int\mathcal{D}\bar{Q}\mathcal{D}Q\mathcal{D}\bar{T}\mathcal{D}T\mathcal{D}G^\alpha_{1\mu}\mathcal{D}G^\alpha_{2\mu}\mathcal{D}B^A_\mu\mathcal{D}Z'_\mu\\ &\quad\times\exp\left(iS_{\mathrm{SBS}}[G^\alpha_{1\mu},G^\alpha_{2\mu},A^A_{1\mu},A^A_{2\mu},W^a_\mu,B_{1\mu},B_{2\mu},\bar{Q},Q,\bar{T},T]\right)\\ &= \mathcal{N}[W^a_\mu,B_\mu]\int\mathcal{D}\mu(U)\exp\left(iS_{\mathrm{eff}}[U,W^a_\mu,B_\mu]\right)\,, \qquad (55)\end{aligned}$$

where $U(x)$ is a dimensionless unitary unimodular matrix field in the electroweak chiral Lagrangian, and $\mathcal{D}\mu(U)$ denotes normalized functional integration measure on U. The normalization factor $\mathcal{N}[W^a_\mu,B_\mu]$ is determined through requirement that when the gauge coupling g_{32} is switched off $S_{\mathrm{eff}}[U,W^a_\mu,B_\mu]$ vanishes, this leads the electroweak gauge fields W^a_μ, B_μ dependent part of $\mathcal{N}[W^a_\mu,B_\mu]$ is

$$\begin{aligned}\mathcal{N}[W^a_\mu,B_\mu] &= \int\mathcal{D}\bar{Q}\mathcal{D}Q\mathcal{D}\bar{T}\mathcal{D}T\mathcal{D}G^\alpha_{1\mu}\mathcal{D}B^A_\mu\mathcal{D}Z'_\mu \qquad (56)\\ &\quad\times\exp\left(iS_{\mathrm{SBS}}[G^\alpha_{1\mu},0,A^A_{1\mu},A^A_{2\mu},W^a_\mu,B_{1\mu},B_{2\mu},\bar{Q},Q,\bar{T},T]\right)\end{aligned}$$

Since there different interactions in present model, in following several subsections, we discuss them and their contributions to EWCL separately.

A. Topcolor Symmetry Breaking: the Contribution of $SU(3)_{\mathrm{TC1}}$

It can be shown below, by Schwinger-Dyson analysis, that the $SU(3)_{\mathrm{TC1}}$ interaction induces the techniquark condensate $\langle\bar{Q}Q\rangle\neq 0$, which will trigger the topcolor symmetry breaking $SU(3)_1\times SU(3)_2\times U(1)_{Y_1}\times U(1)_{Y_2}\to SU(3)_c\times U(1)_Y$ at the scale $\Lambda=1$ TeV. This typically leaves a degenerate, massive color octet of "colorons", B^A_μ, and a singlet heavy Z'_μ in the coset space $[SU(3)_1\times SU(3)_2\times U(1)_{Y_1}\times U(1)_{Y_2}]/[SU(3)_c\times U(1)_Y]$. The gluon A^A_μ and coloron B^A_μ (the SM $U(1)_Y$ field B_μ and the $U(1)'$ field Z'_μ) are defined by orthogonal rotations with mixing angle θ (θ'):

$$\begin{pmatrix}A^A_{1\mu} & A^A_{2\mu}\end{pmatrix}=\begin{pmatrix}B^A_\mu & A^A_\mu\end{pmatrix}\begin{pmatrix}\cos\theta & -\sin\theta\\ \sin\theta & \cos\theta\end{pmatrix}\,, \qquad (57a)$$

$$\begin{pmatrix}B_{1\mu} & B_{2\mu}\end{pmatrix}=\begin{pmatrix}Z'_\mu & B_\mu\end{pmatrix}\begin{pmatrix}\cos\theta' & -\sin\theta'\\ \sin\theta' & \cos\theta'\end{pmatrix}\,, \qquad (57b)$$

which lead to

$$-h_1\frac{\lambda^A}{2}\not{A}_1^A P_L - h_2\frac{\lambda^A}{2}\not{A}_2^A P_R = -g_3\frac{\lambda^A}{2}\not{A}^A - g_3(\cot\theta P_L - \tan\theta P_R)\frac{\lambda^A}{2}\not{B}^A\ , \tag{58a}$$

$$-q_1\frac{1}{2}\not{B}_1 P_L - q_2\frac{1}{2}\not{B}_2 P_R = -g_1\frac{1}{2}\not{B} - g_1(\cot\theta' P_L - \tan\theta' P_R)\frac{1}{2}\not{Z}'\ , \tag{58b}$$

with

$$g_3 \equiv h_1\sin\theta = h_2\cos\theta\ , \tag{59a}$$

$$g_1 \equiv q_1\sin\theta' = q_2\cos\theta'\ . \tag{59b}$$

As a first step , we formally integrate out the $SU(3)_{\rm TC1}$ technigluons $G^\alpha_{1\mu}$ in Eq.(55) by introducing full n-point Green's function of the $G^\alpha_{1\mu}$-field $G^{\alpha_1\ldots\alpha_n}_{\mu_1\ldots\mu_n}$

$$\begin{aligned}
&\exp\bigg(iS_{\rm EW}[W^a_\mu, B_\mu]\bigg)\\
&= \exp\bigg[i\int d^4x(-\frac{1}{4}W^a_{\mu\nu}W^{a\mu\nu})\bigg]\int\mathcal{D}\bar{T}\mathcal{D}T\mathcal{D}G^\alpha_{2\mu}\mathcal{D}B^A_\mu\mathcal{D}Z'_\mu\exp\bigg[i\int d^4x(-\frac{1}{4}F^\alpha_{2\mu\nu}F^{\alpha\mu\nu}_2\\
&\quad -\frac{1}{4}A^A_{1\mu\nu}A^{A\mu\nu}_1 - \frac{1}{4}A^A_{2\mu\nu}A^{A\mu\nu}_2 - \frac{1}{4}B_{1\mu\nu}B^{\mu\nu}_1 - \frac{1}{4}B_{2\mu\nu}B^{\mu\nu}_2) + iS_{\rm TC1}[A^A_{1\mu}, A^A_{2\mu}, B_{1\mu}, B_{2\mu}]\\
&\quad +i\int d^4x\ \bar{T}\big[i\not{\partial} - g_{32}r^\alpha_2\not{G}^\alpha_2 - g_2\frac{\tau^a}{2}\not{W}^aP_L - q_2\frac{1}{6}\not{B}_2P_L - q_2(\frac{1}{6}+\frac{\tau^3}{2})\not{B}_2P_R\big]T\bigg]\ ,
\end{aligned} \tag{60}$$

where

$$\begin{aligned}
&\exp\bigg(iS_{\rm TC1}[A^A_{1\mu}, A^A_{2\mu}, B_{1\mu}, B_{2\mu}]\bigg)\\
&\equiv \int\mathcal{D}\bar{Q}\mathcal{D}Q\exp\bigg[i\int d^4x\bar{Q}\big(i\not{\partial} - h_1\frac{\lambda^A}{2}\not{A}^A_1P_L - h_2\frac{\lambda^A}{2}\not{A}^A_2P_R - q_1\frac{1}{2}\not{B}_1P_L - q_2\frac{1}{2}\not{B}_2P_R\big)Q\\
&\quad +\sum_{n=2}^{\infty}\int d^4x_1\ldots d^4x_n\frac{(-ig_{31})^n}{n!}G^{\alpha_1\ldots\alpha_n}_{\mu_1\ldots\mu_n}(x_1,\ldots,x_n)J^{\mu_1}_{1\alpha_1}(x_1)\ldots J^{\mu_n}_{1\alpha_n}(x_n)\bigg]\ .
\end{aligned} \tag{61}$$

$J^\mu_{1\alpha}(x)\equiv\bar{Q}(x)r^\alpha_1\gamma^\mu Q(x)$ is effective source.

Since the total functional derivative of the integrand in Eq.(61) with respect to $\bar{Q}^\sigma(x)$ is zero, (here and henceforth the suffixes σ and ρ are short notations for Lorentz spinor, techniflavor and technicolor indices,) *i.e.*,

$$\begin{aligned}
0 &= \int\mathcal{D}\bar{Q}\mathcal{D}Q\frac{\delta}{\delta\bar{Q}^\sigma(x)}\exp\bigg[\int d^4x(\bar{Q}I+\bar{I}Q) + i\int d^4x\bar{Q}\big(i\not{\partial} - h_1\frac{\lambda^A}{2}\not{A}^A_1P_L - h_2\frac{\lambda^A}{2}\not{A}^A_2P_R\\
&\quad -q_1\frac{1}{2}\not{B}_1P_L - q_2\frac{1}{2}\not{B}_2P_R\big)Q + \sum_{n=2}^{\infty}\int d^4x_1\ldots d^4x_n\frac{(-ig_{31})^n}{n!}G^{\alpha_1\ldots\alpha_n}_{\mu_1\ldots\mu_n}(x_1,\ldots,x_n)\\
&\quad \times J^{\mu_1}_{1\alpha_1}(x_1)\ldots J^{\mu_n}_{1\alpha_n}(x_n)\bigg]\ ,
\end{aligned} \tag{62}$$

where $I(x)$ and $\bar{I}(x)$ are the external sources for, respectively, $\bar{Q}(x)$ and $Q(x)$, then continue the similar procedure from (18) to (21), by *neglecting higher-point Green's functions* and *taking factorization approximation*, we obtain

$$\begin{aligned} 0 &= \delta_{\sigma\rho}\delta(x-y)+i\bigg[i\not{\partial}_x-h_1\frac{\lambda^A}{2}\not{A}_1^A(x)P_L-h_2\frac{\lambda^A}{2}\not{A}_2^A(x)P_R-q_1\frac{1}{2}\not{B}_1(x)P_L \\ &\quad -q_2\frac{1}{2}\not{B}_2(x)P_R\bigg]_{\sigma\tau}\langle Q^\tau(x)\bar{Q}^\rho(y)\rangle-g_{31}^2\int d^4x_2G^{\alpha_1\alpha_2}_{\mu_1\mu_2}(x,x_2) \\ &\quad \times(r_1^{\alpha_1}\gamma^{\mu_1})_{\sigma\tau}(r_1^{\alpha_2}\gamma^{\mu_2})_{\gamma\delta}\big\langle\bar{Q}^\rho(y)Q^\delta(x_2)\big\rangle\big\langle\bar{Q}^\gamma(x_2)Q^\tau(x)\big\rangle\ , \end{aligned} \tag{63}$$

where $\big\langle\mathcal{O}(x)\big\rangle\equiv\big\langle\!\big\langle\mathcal{O}(x)\big\rangle\!\big\rangle_I/\langle\langle\,1\,\rangle\rangle_I\big|_{I=\bar{I}=0}$ and we have defined the notation $\langle\langle\cdots\rangle\rangle_I$ in this section by

$$\begin{aligned} \big\langle\!\big\langle\mathcal{O}(x)\big\rangle\!\big\rangle_I &\equiv \int\mathcal{D}\bar{Q}\mathcal{D}Q\,\mathcal{O}(x)\,\exp\bigg[\int d^4x(\bar{Q}I+\bar{I}Q)+i\int d^4x\bar{Q}\big(i\not{\partial}-h_1\frac{\lambda^A}{2}\not{A}_1^AP_L \\ &\quad -h_2\frac{\lambda^A}{2}\not{A}_2^AP_R-q_1\frac{1}{2}\not{B}_1P_L-q_2\frac{1}{2}\not{B}_2P_R\big)Q+\sum_{n=2}^{\infty}\int d^4x_1\ldots d^4x_n\frac{(-ig_{31})^n}{n!} \\ &\quad \times G^{\alpha_1\ldots\alpha_n}_{\mu_1\ldots\mu_n}(x_1,\ldots,x_n)J^{\mu_1}_{1\alpha_1}(x_1)\ldots J^{\mu_n}_{1\alpha_n}(x_n)\bigg]\ . \end{aligned} \tag{64}$$

Denote the technifermion propagator $S^{\sigma\rho}(x,x')\equiv\langle Q^\sigma(x)\bar{Q}^\rho(x')\rangle$, Eq.(63) can be written as SDE for techniquark propagator,

$$\begin{aligned} 0 &= S^{-1}_{\sigma\rho}(x,y)+i\bigg[i\not{\partial}_x-h_1\frac{\lambda^A}{2}\not{A}_1^A(x)P_L-h_2\frac{\lambda^A}{2}\not{A}_2^A(x)P_R-q_1\frac{1}{2}\not{B}_1(x)P_L-q_2\frac{1}{2}\not{B}_2(x)P_R\bigg]_{\sigma\rho} \\ &\quad \times\delta(x-y)-g_{31}^2G^{\alpha_1\alpha_2}_{\mu_1\mu_2}(x,y)\bigg[r_1^{\alpha_1}\gamma^{\mu_1}S(x,y)r_1^{\alpha_2}\gamma^{\mu_2}\bigg]_{\sigma\rho}\ . \end{aligned} \tag{65}$$

By defining techniquark self energy Σ as

$$\begin{aligned} i\Sigma_{\sigma\rho}(x,y) &\equiv S^{-1}_{\sigma\rho}(x,y)+i\bigg[i\not{\partial}_x-h_1\frac{\lambda^A}{2}\not{A}_1^A(x)P_L-h_2\frac{\lambda^A}{2}\not{A}_2^A(x)P_R-q_1\frac{1}{2}\not{B}_1(x)P_L \\ &\quad -q_2\frac{1}{2}\not{B}_2(x)P_R\bigg]_{\sigma\rho}\delta(x-y)\ , \end{aligned} \tag{66}$$

the SDE (65) can be written as

$$i\Sigma_{\sigma\rho}(x,y)=g_{31}^2G^{\alpha_1\alpha_2}_{\mu_1\mu_2}(x,y)\bigg[r_1^{\alpha_1}\gamma^{\mu_1}S(x,y)r_1^{\alpha_2}\gamma^{\mu_2}\bigg]_{\sigma\rho}\ . \tag{67}$$

Moreover, from the fact that technigluon propagator is diagonal in the adjoint representation space of $SU(3)_{\rm TC1}$, *i.e.*, $G^{\alpha\beta}_{\mu\nu}(x,y)=\delta^{\alpha\beta}G_{\mu\nu}(x,y)$, and techniquark propagator $\langle Q\bar{Q}\rangle$ is diagonal in the fundamental representation space of $SU(3)_{\rm TC1}$, and also $(r_1^\alpha r_1^\alpha)_{ab} = C_2(3)\delta_{ab}$, Eq.(67) is diagonal in indices a, b and diagonal part becomes

$$i\Sigma^{ij}_{\eta\zeta}(x,y) = C_2(3) g_{31}^2 G_{\mu_1\mu_2}(x,y)[\gamma^{\mu_1} S(x,y)\gamma^{\mu_2}]^{ij}_{\eta\zeta} \ , \tag{68}$$

where $\{i,\ j\}$, and $\{\eta,\ \zeta\}$ are, respectively, techniflavor, and Lorentz spinor indices; and the Casimir operator $C_2(3) = (3^2-1)/(2\times 3) = 4/3$.

1. *The Gap Equation*

We first consider the case of $A^A_{1\mu} = A^A_{2\mu} = B_{1\mu} = B_{2\mu} = 0$. The $SU(3)_{\rm TC1}$ technigluon propagator in Landau gauge is

$$G^{\alpha\beta}_{\mu\nu}(x,y) = \delta^{\alpha\beta}\int \frac{d^4p}{(2\pi)^4} e^{-ip(x-y)} \frac{i}{-p^2[1+\Pi(-p^2)]}\left(g_{\mu\nu} - \frac{p_\mu p_\nu}{p^2}\right) , \tag{69}$$

In the case of $A^A_{1\mu} = A^A_{2\mu} = B_{1\mu} = B_{2\mu} = 0$, the $SU(3)_{\rm TC1}$ techniquark self energy and propagator are respectively

$$\Sigma^{ij}_{\eta\zeta}(x,y) = \int \frac{d^4p}{(2\pi)^4} e^{-ip(x-y)}\, \Sigma^{ij}_{\eta\zeta}(-p^2) \qquad S^{ij}_{\eta\zeta}(x,y) = \int \frac{d^4p}{(2\pi)^4} e^{-ip(x-y)} S^{ij}_{\eta\zeta}(p) \ , \tag{70}$$

with $S^{ij}_{\eta\zeta}(p) = i\{1/(\not p - \Sigma(-p^2))\}^{ij}_{\eta\zeta}$. Substituting above equations into the SDE (68), we have

$$\Sigma^{ij}_{\eta\zeta}(-p^2) = \int \frac{d^4q}{(2\pi)^4} \frac{-C_2(3) g_{31}^2}{(p-q)^2[1+\Pi(-(p-q)^2)]}\left[g_{\mu\nu} - \frac{(p-q)_\mu (p-q)_\nu}{(p-q)^2}\right]\left[\gamma^\mu \frac{i}{\not q - \Sigma(-q^2)}\gamma^\nu\right]^{ij}_{\eta\zeta} \tag{71}$$

As discussions of dynamical computation prescription for one-doublet technicolor model, above equation will lead $\Sigma^{ij}_{\eta\zeta}(-p^2) = \delta^{ij}\delta_{\eta\zeta}\Sigma_{\rm TC}(p_E^2)$ and in Euclidean space $\Sigma_{\rm TC}(p_E^2)$ satisfy

$$\Sigma_{\rm TC1}(p_E^2) = 3C_2(3)\int \frac{d^4q_E}{4\pi^3} \frac{\alpha_{31}[(p_E-q_E)^2]}{(p_E-q_E)^2} \frac{\Sigma_{\rm TC1}(q_E^2)}{q_E^2 + \Sigma^2_{\rm TC1}(q_E^2)} \ . \tag{72}$$

The corresponding techniquark condensate $\langle \bar Q^k Q^j\rangle$ with k and j techniflavor indices,

$$\langle \bar Q^k(x) Q^j(x)\rangle = -12\delta^{jk}\int \frac{d^4p_E}{(2\pi)^4} \frac{\Sigma_{\rm TC}(p_E^2)}{p_E^2 + \Sigma^2_{\rm TC}(p_E^2)} \ , \tag{73}$$

where ${\rm tr}_{lc}$ is the trace with respect to Lorentz, technicolor indices. Nonzero techniquark self energy can give a nontrivial diagonal condensate $\langle \bar Q Q\rangle \neq 0$, which spontaneously breaks $SU(3)_1 \times SU(3)_2 \times U(1)_{Y_1} \times U(1)_{Y_2} \to SU(3)_c \times U(1)_Y$.

In the following we consider the effects of the nonzero electroweak gauge fields $A^A_{1\mu}$, $A^A_{2\mu}$, $B_{1\mu}$ and $B_{2\mu}$. The SDE (68) is explicitly

$$\begin{aligned}\Sigma(x,y) \;=\; & C_2(3)g_{31}^2 G_{\mu\nu}(x,y)\gamma^\mu\Bigg[\bigg(i\not{\partial}_x - h_1\frac{\lambda^A}{2}\not{A}_1^A(x)P_L - h_2\frac{\lambda^A}{2}\not{A}_2^A(x)P_R - q_1\frac{1}{2}\not{B}_1(x)P_L \\ & -q_2\frac{1}{2}\not{B}_2(x)P_R\bigg)\delta(x-y) - \Sigma(x,y)\Bigg]^{-1}\gamma^\nu \;, \qquad (74)\end{aligned}$$

where the techniflavor and Lorentz spinor indices of the techniquark self energy are implicitly contained.

Suppose the function $\Sigma_{\rm TC1}(-p^2)$ is a solution of the SDE in the case $A^A_{1\mu} = A^A_{2\mu} = B_{1\mu} = B_{2\mu} = 0$, that is, it satisfies the equation

$$\Sigma_{\rm TC1}(-p^2) \;=\; C_2(3)g_{31}^2\int\frac{d^4q}{(2\pi)^4}G_{\mu\nu}(q^2)\gamma^\mu\frac{1}{\not{q}+\not{p}-\Sigma_{\rm TC1}[-(q+p)^2]}\gamma^\nu \;, \qquad (75)$$

Replacing the variable p by $p+\Delta$ in Eq.(75) and subsequently integrating over p with the weight $e^{-ip(x-y)}$, we obtain, as long as Δ is commutative with ∂_x and Dirac matrices,

$$\begin{aligned}&\Sigma_{\rm TC1}[(\partial_x - i\Delta)^2]\delta(x-y) \\ &= C_2(3)g_{31}^2 G_{\mu\nu}(x,y)\gamma^\mu\frac{1}{i\not{\partial}_x + \not{\Delta} - \Sigma_{\rm TC1}[-(i\partial_x+\Delta)^2]}\delta(x-y)\gamma^\nu \;. \qquad (76)\end{aligned}$$

Even if Δ is noncommutative with ∂_x and Dirac matrices, the above equation holds as the lowest order approximation, for the commutator $[\not{\partial},\Delta]$ is higher order of momentum than Δ itself. Now if we take Δ to be $-h_1\frac{\lambda^A}{2}A^A_1P_L - h_2\frac{\lambda^A}{2}A^A_2P_R - q_1\frac{1}{2}B_1P_L - q_2\frac{1}{2}B_2P_R$, *ignoring its noncommutative property with ∂_x and Dirac matrices*, Eq.(76) is just the SDE (74) in the case $A^A_{1\mu}\neq 0,\ A^A_{2\mu}\neq 0,\ B_{1\mu}\neq 0,$ and $B_{2\mu}\neq 0$. Thus, $\Sigma_{\rm TC1}[(\partial^x_\mu + ih_1\frac{\lambda^A}{2}A^A_{1\mu}P_L + ih_2\frac{\lambda^A}{2}A^A_{2\mu}P_R + iq_1\frac{1}{2}B_{1\mu}P_L + iq_2\frac{1}{2}B_{2\mu}P_R)^2]\delta(x-y)$, which is $SU(3)_1\times SU(3)_2\times U(1)_{Y_1}\times U(1)_{Y_2}$ covariant, can be regarded as the lowest-order solution of Eq.(74). From Eqs.(58a, 58b), we can write the covariant derivative of $SU(3)_1\times SU(3)_2\times U(1)_{Y_1}\times U(1)_{Y_2}$ as

$$\begin{aligned}\nabla_\mu \;\equiv\; & \partial_\mu + ih_1\frac{\lambda^A}{2}A^A_{1\mu}P_L + ih_2\frac{\lambda^A}{2}A^A_{2\mu}P_R + iq_1\frac{1}{2}B_{1\mu}P_L + iq_2\frac{1}{2}B_{2\mu}P_R \qquad (77)\\ =\; & \partial_\mu + ig_3\frac{\lambda^A}{2}A^A_\mu + ig_1\frac{1}{2}B_\mu + ig_3(\cot\theta P_L - \tan\theta P_R)\frac{\lambda^A}{2}B^A_\mu + ig_1(\cot\theta' P_L - \tan\theta')\frac{1}{2}Z'_\mu \;,\end{aligned}$$

where A^A_μ and B_μ are the gauge fields of the unbroken symmetry group $SU(3)_c\times U(1)_Y$. To further simplify the calculations, we can *just keep this $SU(3)_c\times U(1)_Y$ covariance of the self energy*, that is, we can replace ∇_μ by the covariant derivative of $SU(3)_c\times U(1)_Y$,

$$\overline{\nabla}_\mu \equiv \partial_\mu + ig_3\frac{\lambda^A}{2}A^A_\mu + ig_1\frac{1}{2}B_\mu \qquad (78)$$

inside the techniquark self energy. Thus, if the function $\Sigma(\partial_x^2)\delta(x-y)$ is the self-energy solution of the SDE in the case $A^A_{1\mu}=A^A_{2\mu}=B_{1\mu}=B_{2\mu}=0$, we can replace its argument ∂_x by the $SU(3)_c\times U(1)_Y$ covariant derivative $\overline{\nabla}_x$, *i.e.*, $\Sigma(\overline{\nabla}^2_x)\delta(x-y)$, as an approximate solution the SDE in the case $A^A_{1\mu}\neq 0,\ A^A_{2\mu}\neq 0,\ B_{1\mu}\neq 0,$ and $B_{2\mu}\neq 0$.

Now we are ready to integrate out the techniquarks Q and $\bar{Q}$. The exponential terms on the right-hand side of Eq.(61) can be written explicitly as

$$\begin{aligned}&\sum_{n=2}^{\infty}\int d^4x_1\ldots d^4x_n\frac{(-ig_{31})^n}{n!}G^{\alpha_1\ldots\alpha_n}_{\mu_1\ldots\mu_n}(x_1,\ldots,x_n)J^{\mu_1}_{1,\alpha_1}(x_1)\ldots J^{\mu_n}_{1,\alpha_n}(x_n)\\ &\approx\int d^4xd^4x'\bar{Q}^{\sigma}(x)\Pi_{\sigma\rho}(x,x')Q^{\rho}(x')\ ,\end{aligned}\tag{79}$$

where in the last equality we have taken the approximation of *replacing the summation over* $2n$*-fermion interactions with parts of them by their vacuum expectation values*, that is,

$$\Pi_{\sigma\rho}(x,x')\ =\ \sum_{n=2}^{\infty}\Pi^{(n)}_{\sigma\rho}(x,x')\ ,\tag{80}$$

$$\begin{aligned}\Pi^{(n)}_{\sigma\rho}(x,x')\ &=\ n\times\int d^4x_2\ldots d^4x_{n-1}\frac{(-ig_{31})^n}{n!}G^{\alpha_1\ldots\alpha_n}_{\mu_1\ldots\mu_n}(x,x_2\ldots,x_{n-1},x')\Big\langle(r_1^{\alpha_1}\gamma^{\mu_1})_{\sigma\sigma_1}Q^{\sigma_1}(x)\\ &\times\bar{Q}(x_2)r_1^{\alpha_2}\gamma^{\mu_2}Q(x_2)\ldots\bar{Q}(x_{n-1})r_1^{\alpha_{n-1}}\gamma^{\mu_{n-1}}Q(x_{n-1})\bar{Q}^{\rho_n}(x')(r_1^{\alpha_n}\gamma^{\mu_n})_{\rho_n\rho}\Big\rangle\ ,\end{aligned}\tag{81}$$

where the factor n comes from n different choices of unaveraged $\bar{Q}Q$, and the lowest term of which is

$$\begin{aligned}\Pi^{(2)}_{\sigma\rho}(x,x')\ &=\ 2\cdot\frac{(-ig_{31})^2}{2!}G^{\alpha_1\alpha_2}_{\mu_1\mu_2}(x,x')\Big\langle(r_1^{\alpha_1}\gamma^{\mu_1})_{\sigma\sigma_1}Q^{\sigma_1}(x)\bar{Q}^{\rho_2}(x')(r_1^{\alpha_2}\gamma^{\mu_2})_{\rho_2\rho}\Big\rangle\\ &=\ -g_{31}^2G^{\alpha_1\alpha_2}_{\mu_1\mu_2}(x,x')\Big[r_1^{\alpha_1}\gamma^{\mu_1}S(x,y)r_1^{\alpha_2}\gamma^{\mu_2}\Big]_{\sigma\rho}\ .\end{aligned}\tag{82}$$

Comparing Eq.(82) with Eq.(67), we have

$$i\Pi^{(2)}_{\sigma\rho}(x,x')=\Sigma_{\sigma\rho}(x,x')\approx\Sigma_{\sigma\rho}(\overline{\nabla}^2_x)\delta(x-y)\ .\tag{83}$$

Substituting Eq.(79) into Eq.(61), we obtain

$$\begin{aligned}&\exp\Big(iS_{\rm TC1}[A^A_{1\mu},A^A_{2\mu},B_{1\mu},B_{2\mu}]\Big)\\ &\approx\ \mathrm{Det}\Big[i\partial\!\!\!/-g_3\frac{\lambda^A}{2}A\!\!\!/^A-g_3(\cot\theta P_L-\tan\theta P_R)\frac{\lambda^A}{2}B\!\!\!/^A-g_1\frac{1}{2}B\!\!\!/-g_1(\cot\theta' P_L-\tan\theta' P_R)\frac{1}{2}Z\!\!\!/'\\ &\quad-\Sigma_{\rm TC1}(\overline{\nabla}^2)\Big]\ ,\end{aligned}\tag{84}$$

where we have taken further approximation of *keeping only the lowest order, i.e.* $\Pi^{(2)}_{\sigma\rho}(x,x')$, *of* $\Pi_{\sigma\rho}(x,x')$. With all these approximations, we have

$$\begin{aligned} iS_{\rm TC1}[A^A_\mu,B^A_\mu,B_\mu,Z'_\mu] \;=\; & \mathrm{Tr}\log\bigg[i\not{\partial}-g_3\frac{\lambda^A}{2}\not{A}^A-g_3(\cot\theta P_L-\tan\theta P_R)\frac{\lambda^A}{2}\not{B}^A-g_1\frac{1}{2}\not{B} \\ & -g_1(\cot\theta' P_L-\tan\theta' P_R)\frac{1}{2}\not{Z}'-\Sigma_{\rm TC1}(\overline{\nabla}^2)\bigg]\;, \end{aligned} \tag{85}$$

Since we know QCD-induced condensate is too weak to give sufficiently large masses of W and Z bosons and thus it is negligible when we consider the main cause responsible for the electroweak symmetry breaking which imply that ordinary QCD gluon fields has very little effects on our technicolor and electroweak interactions, therefore for simplicity, we ignore them by just vanishing gluon field $A^A_\mu=0$. In next two subsubsections, we perform low energy expansion and explicitly expand above action up to order of p^4.

2. Low Energy Expansion for $iS_{\rm TC1}[0,B^A_\mu,B_\mu,Z'_\mu]$

We have

$$\begin{aligned} iS_{\rm TC1}[0,B^A_\mu,B_\mu,Z'_\mu] \;&=\; \mathrm{Tr}\log[i\not{\partial}+\not{v}_1+\not{a}_1\gamma_5-\Sigma_{\rm TC1}(\overline{\nabla}^2)] \\ &=\; i\int d^4x(F_0^{\rm TC1})^2\mathrm{tr}[a_1^2(x)]+S^{(4)}_{\rm TC1}[0,B^A_\mu,B_\mu,Z'_\mu]+O(p^6)\;, \end{aligned} \tag{86}$$

where the parameter $F_0^{\rm TC1}$ is depend on the techniquark self energy $\Sigma_{\rm TC1}$. The fields v_μ and a_μ are identified with

$$v_{1,\mu}\equiv-\frac{g_3}{2}(\cot\theta-\tan\theta)\frac{\lambda^A}{2}B^A_\mu-g_1\frac{1}{2}B_\mu-\frac{g_1}{4}(\cot\theta'-\tan\theta')Z'_\mu \tag{87a}$$

$$a_{1,\mu}\equiv-\frac{g_3}{2}(\cot\theta+\tan\theta)\frac{\lambda^A}{2}B^A_\mu-\frac{g_1}{4}(\cot\theta'+\tan\theta')Z'_\mu\;. \tag{87b}$$

Substituting Eqs.(87) into Eq.(86), we obtain, at the order of p^2,

$$\begin{aligned} & S^{(2)}_{\rm TC1}[0,B^A_\mu,B_\mu,Z'_\mu] \\ &=\frac{(F_0^{\rm TC1})^2}{16}\int d^4x[2g_3^2(\cot\theta+\tan\theta)^2B^A_\mu B^{A,,\mu}+3g_1^2(\cot\theta'+\tan\theta')^2Z'^2]\;. \end{aligned} \tag{88}$$

Now we come to consider the p^4 order effective action. It can be divided into two parts

$$S^{(4)}_{\rm TC1}[0,B^A_\mu,B_\mu,Z'_\mu]=S^{(4d)}_{\rm TC1}[0,B^A_\mu,B_\mu,Z'_\mu]+S^{(4c)}_{\rm TC1}[0,B^A_\mu,B_\mu,Z'_\mu] \tag{89}$$

$$iS^{(4d)}_{\rm TC1}[0,B^A_\mu,B_\mu,Z'_\mu]=\mathrm{Tr}\log[i\not{\partial}+\not{v}_1+\not{a}_1\gamma_5] \tag{90}$$

$$iS^{(4c)}_{\rm TC1}[0,B^A_\mu,B_\mu,Z'_\mu]=\mathrm{Tr}\log[i\not{\partial}+\not{v}_1+\not{a}_1\gamma_5-\Sigma_{\rm TC1}(\overline{\nabla}^2)]-\mathrm{Tr}\log[i\not{\partial}+\not{v}_1+\not{a}_1\gamma_5]$$

$S^{(4d)}_{\rm TC1}[0,B^A_\mu,B_\mu,Z'_\mu]$ is divergent part of action, which can calculated by following standard formula

$$iTr\log[i\partial\!\!\!/+l\!\!\!/P_L+r\!\!\!/P_R]=-\frac{1}{2}\mathcal{K}\int d^4x\ {\rm tr}[r^{\mu\nu}r_{\mu\nu}+l^{\mu\nu}l_{\mu\nu}] \tag{91}$$

$$r_{\mu\nu}=\partial_\mu r_\nu-\partial_\nu r_\mu-i(r_\mu r_\nu-r_\nu r_\mu) \qquad\qquad l_{\mu\nu}=\partial_\mu l_\nu-\partial_\nu l_\mu-i(l_\mu l_\nu-l_\nu l_\mu)$$

$$\mathcal{K}=-\frac{1}{48\pi^2}(\log\frac{\kappa^2}{\Lambda^2}+\gamma) \tag{92}$$

with $\mathcal{K}$ a divergent constant depend on the ratio between ultraviolet cutoff Λ and infrared cutoff κ of the theory. Identify

$$r_\mu \ = \ v_{1,\mu}+a_{1,\mu}=-g_3\cot\theta\frac{\lambda^A}{2}B^A_\mu-g_1\frac{1}{2}B_\mu-\frac{g_1}{2}\cot\theta' Z'_\mu \tag{93}$$

$$l_\mu \ = \ v_{1,\mu}-a_{1,\mu}=g_3\tan\theta\frac{\lambda^A}{2}B^A_\mu-g_1\frac{1}{2}B_\mu+\frac{g_1}{2}\tan\theta' Z'_\mu \tag{94}$$

With these preparations,

$$\begin{aligned}S^{(4d)}_{\rm TC1}[0,B^A_\mu,B_\mu,Z'_{c,\mu}] \ = \ & -\frac{1}{2}\mathcal{K}\int d^4x\ \bigg[\frac{g_3^2}{2}(\cot^2\theta B^A_{r,\mu\nu}B^{A,\mu\nu}_r+\tan^2\theta B^A_{l,\mu\nu}B^{A,\mu\nu}_l)+\frac{3g_1^2}{2}B_{\mu\nu}B^{\mu\nu}\\ & +\frac{3g_1^2}{4}(\cot^2\theta'+\tan^2\theta')Z'_{\mu\nu}Z'^{,\mu\nu}+\frac{3g_1^2}{2}(\cot\theta'-\tan\theta')B_{\mu\nu}Z'^{,\mu\nu}\bigg]\end{aligned}$$

with

$$B^A_{r,\mu\nu}=\partial_\mu B^A_\nu-\partial_\nu B^A_\mu-g_3\cot\theta f^{ABC}B^B_\mu B^C_\nu \qquad B^A_{l,\mu\nu}=\partial_\mu B^A_\nu-\partial_\nu B^A_\mu+g_3\tan\theta f^{ABC}B^B_\mu B^C_\nu$$

$$B_{\mu\nu}=\partial_\mu B_\nu-\partial_\nu B_\mu \qquad\qquad Z'_{\mu\nu}=\partial_\mu Z'_\nu-\partial_\nu Z'_\mu \tag{95}$$

$S^{(4c)}_{\rm TC1}[0,B^A_\mu,B_\mu,Z'_\mu]$ is convergent part of action, which can calculated by following standard formula

$$\begin{aligned}S^{(4c)}_{\rm TC1}[0,B^A_\mu,B_\mu,Z'_\mu] \ = \ & \int d^4x\ {\rm tr}[-\mathcal{K}^{{\rm TC1},\Sigma\neq0}_1(d_\mu a^\mu_1)^2-\mathcal{K}^{{\rm TC1},\Sigma\neq0}_2(d_\mu a_{1,\nu}-d_\nu a_{1,\mu})^2+\mathcal{K}^{{\rm TC1},\Sigma\neq0}_3(a^2_1)^2\\ & +\mathcal{K}^{{\rm TC1},\Sigma\neq0}_4(a_{1,\mu}a_{1,\nu})^2-\mathcal{K}^{{\rm TC1},\Sigma\neq0}_{13}V_{1,\mu\nu}V^{\mu\nu}_1+i\mathcal{K}^{{\rm TC1},\Sigma\neq0}_{14}V_{1,\mu\nu}a^\mu_1a^\nu_1]\ ,\end{aligned} \tag{96}$$

with $V_{1,\mu\nu}\equiv\partial_\mu v_{1,\nu}-\partial_\nu v_{1,\mu}-i(v_{1,\mu}v_{1,\nu}-v_{1,\nu}v_{1,\mu})$ and $d_\mu a_{1,\nu}\equiv\partial_\mu a_{1,\nu}-i(v_{1,\mu}a_{1,\nu}-a_{1,\nu}v_{1,\mu})$. $S^{(4c)}_{\rm TC1}[0,B^A_\mu,B_\mu,Z'_\mu]$ can be further divided into four parts

$$\begin{aligned}S^{(4c)}_{\rm TC1}[0,B^A_\mu,B_\mu,Z'_\mu] \ = \ & S^{(4c,B^A)}_{\rm TC1}[B^A]+S^{(4c,B)}_{\rm TC1}[B]+S^{(4c,Z')}_{\rm TC1}[Z']\\ & +S^{(4c,B^AZ')}_{\rm TC1}[B^A,Z']+S^{(4c,BZ')}_{\rm TC1}[B,Z']\end{aligned} \tag{97}$$

The detail form of $S^{(4c,B^A)}_{\rm TC1}[B^A]$, $S^{(4c,B)}_{\rm TC1}[B]$, $S^{(4c,Z')}_{\rm TC1}[Z']$, $S^{(4c,B^AZ')}_{\rm TC1}[B^A,Z']$ and $S^{(4c,BZ')}_{\rm TC1}[B,Z']$ are given in (B1)-(B5) respectively. Since TC1 interaction is $SU(3)$ gauge interaction which is same as QCD interaction and the quark number N_f are all equal to three [1], as we discussed before, due to scale invariance we have $\mathcal{K}_i^{\mathrm{TC1},\Sigma\neq 0}$, $i = 2,3,4,13,14$ are equal to those of QCD values within our approximations

$$\mathcal{K}_i^{\mathrm{TC1},\Sigma\neq 0} = \mathcal{K}_i^{\Sigma\neq 0} \qquad\qquad i = 2,3,4,13,14 \tag{98}$$

Use relation given in Ref.[12]

$$H_1 = -\frac{1}{4}(\mathcal{K}_2^{\Sigma\neq 0} + \mathcal{K}_{13}^{\Sigma\neq 0}) \qquad L_{10} = \frac{1}{2}(\mathcal{K}_2^{\Sigma\neq 0} - \mathcal{K}_{13}^{\Sigma\neq 0}) \qquad L_9 = \frac{1}{8}(4\mathcal{K}_{13}^{\Sigma\neq 0} - \mathcal{K}_{14}^{\Sigma\neq 0}) \tag{99}$$

$$L_1 = \frac{1}{32}(\mathcal{K}_4^{\Sigma\neq 0} + 2\mathcal{K}_{13}^{\Sigma\neq 0} - \mathcal{K}_{14}^{\Sigma\neq 0}) \qquad L_3 = \frac{1}{16}(\mathcal{K}_3^{\Sigma\neq 0} - 2\mathcal{K}_4^{\Sigma\neq 0} - 6\mathcal{K}_{13}^{\Sigma\neq 0} + 3\mathcal{K}_{14}^{\Sigma\neq 0}) \tag{100}$$

In Table III, we list down original QCD calculation result given in Ref.[12], the value for H_1 in the original paper is a divergent constant therefore not given its value, now the divergent part is already extracted out by $S_{\mathrm{TC1}}^{(4d)}[0, B_\mu^A, B_\mu, Z'_\mu]$, H_1 here is a convergent quantity which can be obtained in original formula for H_1 by subtracting out its divergent part caused by terms with $\Sigma = 0$.

TABLE III. The obtained nonzero values of the $O(p^4)$ coefficients $H_1, L_{10}, L_9, L_2, L_1, L_3$ for topcolor-assisted technicolor model.

The F_0^{TC1} and Λ_{TC1} are in units of TeV and coefficients are in units of 10^{-3}.

F_0^{TC1}	Λ_{TC1}	H_1	L_{10}	L_9	L_2	L_1	L_3
1	5.21	43.0	-7.04	5.06	2.19	1.10	-7.81

We finally obtain

$$\mathcal{K}_2^{\mathrm{TC1},\Sigma\neq 0} = L_{10} - 2H_1 \qquad \mathcal{K}_3^{\mathrm{TC1},\Sigma\neq 0} = 64L_1 + 16L_3 + 8L_9 + 2L_{10} + 4H_1$$
$$\mathcal{K}_4^{\mathrm{TC1},\Sigma\neq 0} = 32L_1 - 8L_9 - 2L_{10} - 4H_1$$
$$\mathcal{K}_{13}^{\mathrm{TC1},\Sigma\neq 0} = -L_{10} - 2H_1 \qquad \mathcal{K}_{14}^{\mathrm{TC1},\Sigma\neq 0} = -4L_{10} - 8L_9 - 8H_1 \tag{101}$$

B. Electroweak Symmetry Breaking: the Contribution of $SU(3)_{\mathrm{TC2}}$

Likewise, it is easily check that the $SU(3)_{\mathrm{TC2}}$ interaction does induce the techniquark condensate $\langle \bar{T}T \rangle \neq 0$, which triggers the electroweak symmetry breaking $SU(2)_L \times U(1)_Y \rightarrow$

---

[1] This is in fact an approximation in which we have ignored possible effects on the running of TC1 gauge coupling constant from ordinary color gauge fields and coloron fields.

$U(1)_{\rm EM}$. Integrating out the $SU(3)_{\rm TC2}$ technigluons $G^\alpha_{2\mu}$ and the techniquarks T and $\bar{T}$, Eq.(60) can be written as

$$
\begin{aligned}
&\exp\bigg(iS_{\rm EW}[W^a_\mu,B_\mu]\bigg)\\
&= \exp\bigg[i\int d^4x(-\frac{1}{4}W^a_{\mu\nu}W^{a\mu\nu})\bigg]\int\mathcal{D}B^A_\mu\mathcal{D}Z'_\mu\exp\bigg[i\int d^4x(-\frac{1}{4}A^A_{1\mu\nu}A^{A\mu\nu}_1 \qquad (102)\\
&\quad -\frac{1}{4}A^A_{2\mu\nu}A^{A\mu\nu}_2-\frac{1}{4}B_{1\mu\nu}B^{\mu\nu}_1-\frac{1}{4}B_{2\mu\nu}B^{\mu\nu}_2)+iS_{\rm TC1}[A^A_\mu,B^A_\mu,B_\mu,Z'_\mu]+iS_{\rm TC2}[W^a_\mu,B_{2\mu}]\bigg]\;,
\end{aligned}
$$

where $S_{\rm TC1}[A^A_\mu,B^A_\mu,B_\mu,Z'_\mu]$ has been given in Eq.(85) for its general form and expanded up to order of p^2 in (88) and p^4 in (89), and $S_{\rm TC2}[W^a_\mu,B_{2\mu}]$ is given by

$$
\begin{aligned}
&\exp\bigg(iS_{\rm TC2}[W^a_\mu,B_{2\mu}]\bigg) \qquad (103)\\
&=\int\mathcal{D}\bar{T}\mathcal{D}T\mathcal{D}G^\alpha_{2\mu}\exp\bigg[i\int d^4x\bigg(-\frac{1}{4}F^\alpha_{2\mu\nu}F^{\alpha\mu\nu}_2+\bar{T}\big[i\partial\!\!\!/-g_{32}r^\alpha_2 G\!\!\!/^\alpha_2+l\!\!\!/_2P_L+r\!\!\!/_2P_R\big]T\bigg)\bigg]
\end{aligned}
$$

with $l_{2,\mu}\equiv-g_2\frac{\tau^a}{2}W^a_\mu-q_2\frac{1}{6}B_{2\mu}$ and $r_{2,\mu}\equiv-q_2(\frac{1}{6}+\frac{\tau^3}{2})B_{2\mu}$.

By means of the Gasser-Leutwyler's prescription presented in section II, the functional integration (103) can be related to the QCD-type chiral Lagrangian by

$$
\begin{aligned}
&\frac{\int\mathcal{D}\bar{T}\mathcal{D}T\mathcal{D}G^\alpha_{2\mu}\exp\bigg[i\int d^4x\bigg(-\frac{1}{4}F^\alpha_{2\mu\nu}F^{\alpha\mu\nu}_2+\bar{T}\big[i\partial\!\!\!/-g_{32}r^\alpha_2 G\!\!\!/^\alpha_2+l\!\!\!/_2P_L+r\!\!\!/_2P_R\big]T\bigg)\bigg]}{\int\mathcal{D}\bar{T}\mathcal{D}T\exp\big\{i\int d^4x\bar{T}[i\partial\!\!\!/+l\!\!\!/_2P_L+r\!\!\!/_2P_R]T\big\}}\\
&=\int\mathcal{D}\mu(\tilde{U})\exp\{iS_{\text{TC2-induced eff}}[\tilde{U},l_{2,\mu},r_{2,\mu}]\}\;, \qquad (104)
\end{aligned}
$$

with the $SU(3)_{\rm TC2}$-induced chiral effective action

$$
\begin{aligned}
S_{\text{TC2-induced eff}}[\tilde{U},l_{2,\mu},r_{2,\mu}] \;=\; &\int d^4x\bigg[\frac{(F^{\rm TC2}_0)^2}{4}{\rm tr}[(\nabla^\mu\tilde{U}^\dagger)(\nabla_\mu\tilde{U})]+L^{\rm TC2}_1[{\rm tr}(\nabla^\mu\tilde{U}^\dagger\nabla_\mu\tilde{U})]^2\\
&+L^{\rm TC2}_2{\rm tr}[\nabla_\mu\tilde{U}^\dagger\nabla_\nu\tilde{U}]{\rm tr}[\nabla^\mu\tilde{U}^\dagger\nabla^\nu\tilde{U}]+L^{\rm TC2}_3{\rm tr}[(\nabla^\mu\tilde{U}^\dagger\nabla_\mu\tilde{U})^2]\\
&-iL^{\rm TC2}_9{\rm tr}[F^R_{\mu\nu}\nabla^\mu\tilde{U}\nabla^\nu\tilde{U}^\dagger+F^L_{\mu\nu}\nabla^\mu\tilde{U}^\dagger\nabla^\nu\tilde{U}]+L^{\rm TC2}_{10}{\rm tr}[\tilde{U}^\dagger F^R_{\mu\nu}\tilde{U}F^{L,\mu\nu}]\\
&+H^{\rm TC2}_1{\rm tr}[F^R_{\mu\nu}F^{R,\mu\nu}+F^L_{\mu\nu}F^{L,\mu\nu}]\bigg]\;, \qquad (105)
\end{aligned}
$$

where

$$
\begin{aligned}
&\nabla_\mu\tilde{U}\equiv\partial_\mu\tilde{U}-ir_{2,\mu}\tilde{U}+i\tilde{U}l_{2,\mu}\;,\qquad \nabla_\mu\tilde{U}^\dagger=-\tilde{U}^\dagger(\nabla_\mu\tilde{U})\tilde{U}^\dagger=\partial_\mu\tilde{U}^\dagger-il_{2,\mu}\tilde{U}^\dagger+i\tilde{U}^\dagger r_{2,\mu}\;,\\
&F^R_{\mu\nu}\equiv i[\partial_\mu-ir_{2,\mu},\partial_\nu-ir_{2,\nu}]\;,\qquad F^L_{\mu\nu}\equiv i[\partial_\mu-il_{2,\mu},\partial_\nu-il_{2,\nu}]\;. \qquad (106)
\end{aligned}
$$

The coefficients $F_0^{\rm TC2}$, $L_1^{\rm TC2}$, $L_2^{\rm TC2}$, $L_3^{\rm TC2}$, $L_{10}^{\rm TC2}$, $H_1^{\rm TC2}$ arise from $SU(3)_{\rm TC2}$ dynamics. These coefficients relates to the $\mathcal{K}_i^{\rm TC2}$ coefficients as that appeared in one doublet technicolor model as

$$\begin{aligned}
H_1^{\rm TC2} &= -\frac{\mathcal{K}_2^{{\rm TC2},\Sigma'}+\mathcal{K}_{13}^{{\rm TC2},\Sigma'}}{4} \qquad L_{10}^{\rm TC2}=\frac{\mathcal{K}_2^{{\rm TC2},\Sigma\neq0}-\mathcal{K}_{13}^{{\rm TC2},\Sigma\neq0}}{2}\\
L_9^{\rm TC2} &= \frac{\mathcal{K}_{13}^{{\rm TC2},\Sigma\neq0}}{2}-\frac{\mathcal{K}_{14}^{{\rm TC2},\Sigma\neq0}}{8} \qquad L_2^{\rm TC2}=\frac{\mathcal{K}_4^{{\rm TC2},\Sigma\neq0}+2\mathcal{K}_{13}^{{\rm TC2},\Sigma\neq0}-\mathcal{K}_{14}^{{\rm TC2},\Sigma\neq0}}{16}\\
L_1^{\rm TC2}+\frac{L_3^{\rm TC2}}{2} &= \frac{\mathcal{K}_3^{{\rm TC2},\Sigma\neq0}-\mathcal{K}_4^{{\rm TC2},\Sigma\neq0}-4\mathcal{K}_{13}^{{\rm TC2},\Sigma\neq0}+2\mathcal{K}_{14}^{{\rm TC2},\Sigma\neq0}}{32}
\end{aligned} \tag{107}$$

$\mathcal{K}_i^{\rm TC2}$ coefficients with superscript TC2 denote present TC2 interaction, they are functions of technifermion T self energy $\Sigma_{\rm TC2}(p^2)$ and detail expressions are already written down in (36) of Ref.[12] with the replacement of $N_c \rightarrow 3$ and subtract out their $\Sigma_{\rm TC2}(p^2)=0$ parts. Since TC2 interactions among techiquark doublet T is $SU(3)$ which is same as one doublet technicolor model discussed before except $\Lambda_{\rm TC2}$ may be different as $\Lambda_{\rm TC}$ of one doublet technicolor model, but as we discussed before $\mathcal{K}_i^{\rm TC2},\;\; i=1,2,3,4,13,14$ are independent of $\Lambda_{\rm TC2}$, therefore our $\mathcal{K}_i^{\rm TC2},\;\; i=1,2,3,4,13,14$ are same as those obtained in one doublet technicolor model. This results present $L_i^{\rm TC2},\;\; i=1,2,3,9,10$ and $H_1^{\rm TC2}$ coefficients are same as those in one doublet technicolor model,in Table VI, we list down the numerical calculation result in which the method is already mentioned in previous section and except the result for $H_1^{\rm TC2}$, all others are already used in Table I.

TABLE VI. The obtained nonzero values of the $O(p^4)$ coefficients $H_1^{\rm TC2}, L_{10}^{\rm TC2}, L_9^{\rm TC2}, L_2^{\rm TC2}, L_1^{\rm TC2}, L_3^{\rm TC2}$ for topcolor-assisted technicolor model. The $F_0^{\rm TC2}$ and Λ_{TC2} are in units of TeV and coefficients are in units of 10^{-3}.

$F_0^{\rm TC2}$	$\Lambda_{\rm TC2}$	$H_1^{\rm TC2}=H_1^{\rm 1D}$	$L_{10}^{\rm TC2}=L_{10}^{\rm 1D}$	$L_9^{\rm TC2}=L_9^{\rm 1D}$	$L_2^{\rm TC2}=L_2^{\rm 1D}$	$L_1^{\rm TC2}=L_1^{\rm 1D}$	$L_3^{\rm TC2}=L_3^{\rm 1D}$
0.25	1.34	43.0	-6.90	4.87	2.02	1.01	-7.40

Similar as one-doublet case for $\tilde{U}$ is a 2×2 unitary matrix, and thus the $L_1^{\rm TC2}$ term and the $L_3^{\rm TC2}$ term are linearly related,

$$L_3^{\rm TC2}{\rm tr}[(\nabla^\mu\tilde{U}^\dagger\nabla_\mu\tilde{U})^2] \;=\; \frac{L_3^{\rm TC2}}{2}\Big\{{\rm tr}\big[\tilde{U}^\dagger(\nabla^\mu\tilde{U})\tilde{U}^\dagger(\nabla^\mu\tilde{U})\big]\Big\}^2 \tag{108}$$

Comparing Eqs.(106) with standard covariant derivative given in Ref.[2], we need to recognize

$$\tilde{U}^\dagger = U\;, \qquad \nabla_\mu\tilde{U}^\dagger=\tilde{D}_\mu U\equiv\partial_\mu U+ig_2\frac{\tau^a}{2}W_\mu^aU-Uiq_2\frac{\tau^3}{2}B_{2\mu}\;. \tag{109}$$

And $F^R_{\mu\nu}=-q_2(\frac{1}{6}+\frac{\tau^3}{2})B_{2\mu\nu}$, $F^L_{\mu\nu}=-g_2\frac{\tau^a}{2}W^a_{\mu\nu}-q_2\frac{1}{6}B_{2\mu\nu}$ with $B_{2\mu\nu}\equiv\partial_\mu B_{2\nu}-\partial_\nu B_{2\mu}$ is the $U(1)_{Y_2}$ gauge field strength tensor.

Substituting above equations back into Eq.(105), we obtain

$$\begin{aligned}S_{\text{TC2-induced eff}}[U,W,B_2] \;=\; &\int d^4x\bigg[-\frac{(F_0^{\rm TC2})^2}{4}\text{tr}(\tilde{X}_\mu\tilde{X}^\mu)+(L_1^{\rm 1D}+\frac{L_3^{\rm 1D}}{2})[\text{tr}(\tilde{X}_\mu\tilde{X}^\mu)]^2\\ &+L_2^{\rm 1D}[\text{tr}(\tilde{X}_\mu\tilde{X}_\nu)]^2-i\frac{L_9^{\rm 1D}}{2}q_2B_{2\mu\nu}\text{tr}(\tau^3\tilde{X}^\mu\tilde{X}^\nu)-iL_9^{\rm 1D}\text{tr}(\overline{W}_{\mu\nu}\tilde{X}^\mu\tilde{X}^\nu)\\ &+\frac{L_{10}^{\rm 1D}}{2}q_2B_{2\mu\nu}\text{tr}(\tau^3\overline{W}^{\mu\nu})+\frac{1}{18}(L_{10}^{\rm 1D}+11H_1^{\rm 1D})q_2^2B_{2\mu\nu}B_2^{\mu\nu}\\ &+H_1^{\rm 1D}\text{tr}(\overline{W}_{\mu\nu}\overline{W}^{\mu\nu})\bigg]\,, \end{aligned}\tag{110}$$

where $\tilde{X}_\mu$ is defined by

$$\tilde{X}_\mu\equiv U^\dagger(\tilde{D}_\mu U)\,. \tag{111}$$

With (111) and from Eqs.(109), (57b) and (59b), we obtain

$$\tilde{X}_\mu=X_\mu+ig_1\tan\theta' Z'_\mu\frac{\tau^3}{2}\,. \tag{112}$$

Substituting Eq.(112) into Eq.(110), we obtain, at the order of p^2,

$$\begin{aligned}&S^{(2)}_{\text{TC2-induced eff}}[U,W^a,B\cos\theta'-Z'\sin\theta']\\ &=\frac{(F_0^{\rm TC2})^2}{4}\int d^4x\bigg[-\text{tr}(X_\mu X^\mu)-ig_1\tan\theta' Z'_\mu\text{tr}(\tau^3X^\mu)+\frac{g_1^2}{2}\tan^2\theta' Z'^2\bigg]\,.\end{aligned}\tag{113}$$

Similarly detail algebra gives

$$
\begin{aligned}
&iS^{(4)}_{\text{TC2-induced eff}}[U, W^a, B\cos\theta' - Z'\sin\theta'] \\
&= i\int d^4x\bigg[(L_1^{1\mathrm{D}} + \frac{L_3^{1\mathrm{D}}}{2})[\mathrm{tr}(X_\mu X^\mu)]^2 + L_2^{1\mathrm{D}}[\mathrm{tr}(X_\mu X_\nu)]^2 - iL_9^{1\mathrm{D}}\mathrm{tr}(\overline{W}_{\mu\nu}X^\mu X^\nu) \\
&\quad -i\frac{L_9^{1\mathrm{D}}}{2}g_1B_{\mu\nu}\mathrm{tr}(\tau^3X^\mu X^\nu) + \frac{L_{10}^{1\mathrm{D}}}{2}g_1B_{\mu\nu}\mathrm{tr}(\tau^3\overline{W}^{\mu\nu}) + \frac{1}{18}(L_{10}^{1\mathrm{D}} + 11H_1^{1\mathrm{D}})g_1^2B_{\mu\nu}B^{\mu\nu} \\
&\quad +H_1^{1\mathrm{D}}\mathrm{tr}(\overline{W}_{\mu\nu}\overline{W}^{\mu\nu}) - (L_1^{1\mathrm{D}} + \frac{L_3^{1\mathrm{D}}}{2})g_1^2\tan^2\theta' Z'^{,2}(\mathrm{tr}X^2) \qquad (114) \\
&\quad +(L_1^{1\mathrm{D}} + L_2^{1\mathrm{D}} + \frac{L_3^{1\mathrm{D}}}{2})[\frac{1}{4}g_1^4\tan^4\theta' Z'^{,4} - ig_1^3\tan^3\theta' Z'^{,2}Z'^{,\mu}\mathrm{tr}(X_\mu\tau^3)] \\
&\quad -\frac{1}{2}L_2^{1\mathrm{D}}g_1^2\tan^2\theta'[Z'^2\mathrm{tr}(X_\nu\tau^3)\mathrm{tr}(X^\nu\tau^3) + Z'^\mu Z'^\nu\mathrm{tr}(X_\mu\tau^3)\mathrm{tr}(X_\nu\tau^3)] \\
&\quad -(L_1^{1\mathrm{D}} + \frac{L_3^{1\mathrm{D}}}{2})g_1^2\tan^2\theta' Z'^{,\mu}Z'^{,\nu}\mathrm{tr}(X_\mu\tau^3)(X_\nu\tau^3) - L_2^{1\mathrm{D}}g_1^2\tan^2\theta' Z'^{,\mu}Z'^{,\nu}\mathrm{tr}(X_\mu X_\nu) \\
&\quad +2i(L_1^{1\mathrm{D}} + \frac{L_3^{1\mathrm{D}}}{2})g_1\tan\theta' Z'_\mu\mathrm{tr}(X^\mu\tau^3)(\mathrm{tr}X^2) + 2iL_2^{1\mathrm{D}}g_1\tan\theta' Z'_\mu\mathrm{tr}(X_\nu\tau^3)\mathrm{tr}(X^\mu X^\nu) \\
&\quad +i\frac{L_9^{1\mathrm{D}}}{2}g_1\tan\theta' Z'_{\mu\nu}\mathrm{tr}(\tau^3X^\mu X^\nu) + \frac{1}{2}g_1L_9^{1\mathrm{D}}\tan\theta'[\mathrm{tr}(\overline{W}_{\mu\nu}X^\mu\tau^3)Z'_\nu + \mathrm{tr}(\overline{W}_{\mu\nu}\tau^3X^\nu)Z'_\mu] \\
&\quad -\frac{L_{10}^{1\mathrm{D}}}{2}g_1\tan\theta' Z'_{\mu\nu}\mathrm{tr}(\tau^3\overline{W}^{\mu\nu}) + \frac{1}{18}(L_{10}^{1\mathrm{D}} + 11H_1^{1\mathrm{D}})[g_1^2\tan^2\theta' Z'_{\mu\nu}Z'^{,\mu\nu} - 2g_1^2\tan\theta' Z'_{\mu\nu}B^{\mu\nu}]\bigg] .
\end{aligned}
$$

Thus, from Eqs.(103–111) we obtain

$$
\begin{aligned}
iS_{\mathrm{TC2}}[W_\mu^a, B_{2\mu}] &= \mathrm{Tr}\ln\big[i\partial\!\!\!/ - g_2\frac{\tau^a}{2}W\!\!\!\!/^a P_L - q_2\frac{1}{6}B\!\!\!/_2P_L - q_2(\frac{1}{6} + \frac{\tau^3}{2})B\!\!\!/_2P_R\big] \qquad (115) \\
&\quad \times\log\int\mathcal{D}\mu(U)\exp\left(iS^{(2)}_{\text{TC2-induced eff}}[U, W, B_2] + iS^{(4)}_{\text{TC2-induced eff}}[U, W, B_2]\right) .
\end{aligned}
$$

We still left to compute $\mathrm{Tr}\ln\big[i\partial\!\!\!/ - g_2\frac{\tau^a}{2}W\!\!\!\!/^a P_L - q_2\frac{1}{6}B\!\!\!/_2P_L - q_2(\frac{1}{6} + \frac{\tau^3}{2})B\!\!\!/_2P_R\big]$, which is at least order of p^4. We can write $\mathrm{Tr}\ln\big[i\partial\!\!\!/ - g_2\frac{\tau^a}{2}W\!\!\!\!/^a P_L - q_2\frac{1}{6}B\!\!\!/_2P_L - q_2(\frac{1}{6} + \frac{\tau^3}{2})B\!\!\!/_2P_R\big]$ as

$$
\begin{aligned}
&\mathrm{Tr}\ln\big[i\partial\!\!\!/ - g_2\frac{\tau^a}{2}W\!\!\!\!/^a P_L - q_2\frac{1}{6}B\!\!\!/_2P_L - q_2(\frac{1}{6} + \frac{\tau^3}{2})B\!\!\!/_2P_R\big] = \mathrm{Tr}\log[i\partial\!\!\!/ + l\!\!\!/_2P_L + r\!\!\!/_2P_R] \qquad (116) \\
&l_2^\mu = -g_2\frac{\tau^a}{2}W^{a,\mu} - q_2\frac{1}{6}B_2^\mu = -g_2\frac{\tau^a}{2}W^{a,\mu} - \frac{1}{6}(g_1B^\mu - g_1\tan\theta' Z'^{,\mu}) \\
&r_2^\mu = -q_2(\frac{1}{6} + \frac{\tau^3}{2})B_2^\mu = -(\frac{1}{6} + \frac{\tau^3}{2})(g_1B^\mu - g_1\tan\theta' Z'^{,\mu})
\end{aligned}
$$

Then with help of (91), computation gives

$$
\begin{aligned}
&i\mathrm{Tr}\ln\big[i\partial\!\!\!/ - g_2\frac{\tau^a}{2}W\!\!\!\!/^a P_L - q_2\frac{1}{6}B\!\!\!/_2P_L - q_2(\frac{1}{6} + \frac{\tau^3}{2})B\!\!\!/_2P_R\big] \qquad (117) \\
&= -\frac{1}{2}\mathcal{K}\int d^4x\left[\frac{11}{18}g_1^2(B_{\mu\nu}B^{\mu\nu} + \tan^2\theta' Z'_{\mu\nu}Z'^{,\mu\nu} - 2\tan\theta' Z'_{\mu\nu}B^{\mu\nu}) + \frac{1}{2}g_2^2W^a_{\mu\nu}W^{a,\mu\nu}\right]
\end{aligned}
$$

Substituting Eq.(115) into Eq.(102) and then comparing it with the last line of Eq.(55), we have

$$\begin{aligned}
&\mathcal{N}[W^a_\mu, B_\mu]\exp\left(iS_{\text{eff}}[U, W^a_\mu, B_\mu]\right)\\
&= \exp\left[i\int d^4x(-\frac{1}{4}W^a_{\mu\nu}W^{a\mu\nu})\right]\int \mathcal{D}B^A_\mu\mathcal{D}Z'_\mu \exp\left[i\int d^4x(-\frac{1}{4}A^A_{1\mu\nu}A_1^{A\mu\nu} - \frac{1}{4}A^A_{2\mu\nu}A_2^{A\mu\nu}\right.\\
&\quad -\frac{1}{4}B_{1\mu\nu}B_1^{\mu\nu} - \frac{1}{4}B_{2\mu\nu}B_2^{\mu\nu}) + \mathrm{Tr}\ln\left[i\not{\partial} - g_2\frac{\tau^a}{2}\not{W}^a P_L - q_2\frac{1}{6}\not{B}_2 P_L - q_2(\frac{1}{6}+\frac{\tau^3}{2})\not{B}_2 P_R\right]\\
&\quad \left. +iS_{\text{TC1}}[A^A_\mu, B^A_\mu, B_\mu, Z'_\mu] + iS_{\text{TC2-induced eff}}[U, W, B_2]\right]_{A^A_\mu = 0}, \qquad (118)
\end{aligned}$$

where we have put the $SU(3)_c$ gluon fields $A^A_\mu = 0$ on the right-hand side, for the QCD effects are small here. The normalization factor from its definition (56) can be calculated similarly as previous procedure, the only difference is that we switch off TC2 interaction by taking $g_{32} = 0$ and it will result $S_{\text{TC2-induced eff}}[U, W, B_2]$ vanishes, then this leads ignoring term $iS_{\text{TC2-induced eff}}[U, W, B_2]$ in above expression, we get expression for $\mathcal{N}[W^a_\mu, B_\mu]$.

C. Integrate out of Colorons

Now, as shown in Eqs.(118), the next work is to integrate out the $SU(3)_c$ octet of colorons, B^A_μ. From Eqs.(57a) and (59a), it is straightforward to get

$$A^A_{1\mu\nu} = A^A_{\mu\nu}\sin\theta + B^A_{1,\mu\nu}\cos\theta\ , \qquad (119a)$$

$$A^A_{2\mu\nu} = A^A_{\mu\nu}\cos\theta - B^A_{2,\mu\nu}\sin\theta\ , \qquad (119b)$$

where

$$B^A_{1,\mu\nu} \equiv \partial_\mu B^A_\nu - \partial_\nu B^A_\mu + g_3 f^{ABC}(\cot\theta B^B_\mu B^C_\nu + B^B_\mu A^C_\nu + A^B_\mu B^C_\nu) \qquad (120)$$

$$B^A_{2,\mu\nu} \equiv \partial_\mu B^A_\nu - \partial_\nu B^A_\mu - g_3 f^{ABC}(\tan\theta B^B_\mu B^C_\nu + B^B_\mu A^C_\nu + A^B_\mu B^C_\nu) \qquad (121)$$

then Eqs.(118) become

$$\begin{aligned}
&\mathcal{N}[W^a_\mu, B_\mu]\exp\left(iS_{\text{eff}}[U, W^a_\mu, B_\mu]\right)\\
&= \exp\left[i\int d^4x(-\frac{1}{4}W^a_{\mu\nu}W^{a\mu\nu})\right]\int \mathcal{D}B^A_\mu\mathcal{D}Z'_\mu \exp\left[iS_{\text{TC1}}[0, B^A_\mu, B_\mu, Z'_\mu]\right. \qquad (122)\\
&\quad +i\int d^4x(-\frac{1}{4}B^A_{1\mu\nu}B_1^{A\mu\nu}\cos^2\theta - \frac{1}{4}B^A_{2\mu\nu}B_2^{A\mu\nu}\sin^2\theta - \frac{1}{4}B_{1\mu\nu}B_1^{\mu\nu} - \frac{1}{4}B_{2\mu\nu}B_2^{\mu\nu})\\
&\quad \left.+\mathrm{Tr}\ln\left[i\not{\partial} - g_2\frac{\tau^a}{2}\not{W}^a P_L - q_2\frac{1}{6}\not{B}_2 P_L - q_2(\frac{1}{6}+\frac{\tau^3}{2})\not{B}_2 P_R\right] + iS_{\text{TC2-induced eff}}[U, W, B_2]\right],
\end{aligned}$$

Ignoring term $iS_{\text{TC2-induced eff}}[U,W,B_2]$ in above expression, we get expression for $\mathcal{N}[W_\mu^a,B_\mu]$. In above result, if we denote the coloron involved part be $\int\mathcal{D}B_\mu^A\ e^{iS_{\text{coloron}}[B^A,Z']}$, then

$$\begin{aligned}
S_{\text{coloron}}[B^A,Z'] &= S_{\text{TC1}}^{4c,B^A}[B^A]+S_{\text{TC1}}^{4c,B^AZ'}[B^A,Z']+\int d^4x\bigg(-\frac{1}{4}B^A_{1\mu\nu}B_1^{A\mu\nu}\cos^2\theta \\
&\quad -\frac{1}{4}B^A_{2\mu\nu}B_2^{A\mu\nu}\sin^2\theta+\frac{(F_0^{\text{TC1}})^2}{8}g_3^2(\cot\theta+\tan\theta)^2B_\mu^AB^{A,,\mu} \\
&\quad -\frac{g_3^2}{4}\mathcal{K}(\cot^2\theta B^A_{r,\mu\nu}B_r^{A,\mu\nu}+\tan^2\theta B^A_{l,\mu\nu}B_l^{A,\mu\nu})\bigg) \qquad (123)\\
&= S^0_{\text{coloron}}[B^A,Z']+S^{\text{int}}_{\text{coloron}}[B^A,Z'] \qquad (124)
\end{aligned}$$

with $S^0_{\text{coloron}}[B^A,Z']$ be linear and quadratic in coloron fields and $S^{\text{int}}_{\text{coloron}}[B^A,Z']$ be cubic and quartic in coloron fields, the detail form of them are given in (C1) and (C2). Now coloron fields is not correctly normalized, since the coefficient in front of kinetic term is not standard $-1/4$. We now introduce normalized fields $B^A_{R,\mu}$ as

$$B_\mu^A=\frac{1}{c}B^A_{R,\mu} \qquad (125)$$

$$c^2=1+g_3^2[\frac{1}{2}\mathcal{K}_2^{\text{TC1},\Sigma\neq0}(\cot\theta+\tan\theta)^2+\frac{1}{2}\mathcal{K}_{13}^{\text{TC1},\Sigma\neq0}(\cot\theta-\tan\theta)^2+\mathcal{K}(\cot^2\theta+\tan^2\theta)] \qquad (126)$$

With them, $S^0_{\text{coloron}}[B^A,Z']$ in terms of normalized coloron fields become

$$S^0_{\text{coloron}}[B^A,Z']=\int d^4x\ \frac{1}{2}B^A_{R,\mu}(x)D_B^{-1,\mu\nu}(Z')B^A_{R,\nu}(x) \qquad (127)$$

with

$$D_B^{-1,\mu\nu}(Z')=D_{B0}^{-1,\mu\nu}+\Delta^{\mu\nu}(Z') \qquad D_{B0}^{-1,\mu\nu}=g^{\mu\nu}(\partial^2+M^2_{\text{coloron}})-(1+\lambda_B)\partial^\mu\partial^\nu \qquad (128)$$

$$\begin{aligned}
\Delta^{\mu\nu}(Z') &= [g^{\mu\nu}(\frac{1}{2}\mathcal{K}_3^{\text{TC1},\Sigma\neq0}+\mathcal{K}_4^{\text{TC1},\Sigma\neq0})Z'_{\nu'}Z'^{,\nu'}+(2\mathcal{K}_3^{\text{TC1},\Sigma\neq0}+\frac{5}{4}\mathcal{K}_4^{\text{TC1},\Sigma\neq0})Z'^{,\mu}Z'^{,\nu}] \\
&\quad \times\frac{g_1^2g_3^4(\cot\theta+\tan\theta)^2(\cot\theta'+\tan\theta')^2}{32c^2} \qquad (129)
\end{aligned}$$

$$M_{\text{coloron}}=\frac{1}{2}g_3\frac{\cot\theta+\tan\theta}{c}F_0^{\text{TC1}}=\frac{g_3F_0^{\text{TC1}}}{2c\sin\theta\cos\theta} \qquad (130)$$

$$\lambda_B=-\frac{1}{4}g_3^2\frac{(\cot\theta+\tan\theta)^2}{c^2}\mathcal{K}_1^{\text{TC1},\Sigma\neq0} \qquad (131)$$

Here we recover the estimation for coloron mass $M_{\text{coloron}}\sim g_3\Lambda/(\sin\theta\cos\theta)$ given in Ref.[17] if we identify $\Lambda=F_0^{\text{TC1}}/(2c)$. We now denote the result action after integration over colorons as

$$\int\mathcal{D}B_\mu^A\ e^{iS_{\text{coloron}}[B^A,Z']}=e^{i\bar{S}_{\text{coloron}}[Z']} \qquad (132)$$

$\bar{S}_{\rm coloron}[Z']$ are all vacuum diagrams with propagator $D_B^{\mu\nu}(Z')$ and vertices determined by $S^{\rm int}_{\rm coloron}[B^A, Z']$. The loop expansion result is

$$i\bar{S}_{\rm coloron}[Z'] = -\frac{1}{2}\mathrm{Tr}\log D_B^{-1}(Z') + \text{two or more loop contributions} \tag{133}$$

The first term in the r.h.s. of above equation is one loop result, if we further perform low energy expansion for it and drop out total derivative terms, we find the contributions from one loop term is quartically divergent up to order of p^4 which will be vanish if we take dimensional regularization, then up to order of one loop precision, colorons makes no contributions.

D. Integrate out of Z'

From Eqs.(57b) and (59b) imply

$$B_{1\mu\nu} = B_{\mu\nu}\sin\theta' + (\partial_\mu Z'_\nu - \partial_\nu Z'_\mu)\cos\theta' \ , \tag{134a}$$

$$B_{2\mu\nu} = B_{\mu\nu}\cos\theta' - (\partial_\mu Z'_\nu - \partial_\nu Z'_\mu)\sin\theta' \ . \tag{134b}$$

Substituting $B_{1\mu} = B_\mu\sin\theta' + Z'_\mu\cos\theta'$, $B_{2\mu} = B_\mu\cos\theta' - Z'_\mu\sin\theta'$ and Eq.(134) into the right-hand side of Eq.(122), combined with (132), we get

$$\begin{aligned}
&\mathcal{N}[W^a_\mu, B_\mu]\exp\Big(iS_{\rm eff}[U, W^a_\mu, B_\mu]\Big)\\
&= \exp\Big[i\int d^4x(-\frac{1}{4}W^a_{\mu\nu}W^{a\mu\nu})\Big]\int\mathcal{D}Z'_\mu\exp\Big[i\int d^4x\Big(-\frac{1}{4}B_{\mu\nu}B^{\mu\nu} - \frac{1}{4}(\partial_\mu Z'_\nu - \partial_\nu Z'_\mu)^2\\
&\quad+\frac{3g_1^2(F_0^{\rm TC1})^2}{16}(\cot\theta' + \tan\theta')^2 Z'^2 - \frac{1}{2}\mathcal{K}[\frac{3g_1^2}{2}B_{\mu\nu}B^{\mu\nu} + \frac{3g_1^2}{4}(\cot^2\theta' + \tan^2\theta')Z'_{\mu\nu}Z'^{,\mu\nu}\\
&\quad+\frac{3g_1^2}{2}(\cot\theta' - \tan\theta')B_{\mu\nu}Z'^{,\mu\nu}]\Big) + i\bar{S}_{\rm coloron}[Z'] + iS^{(4c,B)}_{\rm TC1}[B] + iS^{(4c,Z')}_{\rm TC1}[Z']\\
&\quad+iS^{(4c,BZ')}_{\rm TC1}[B, Z'] + \mathrm{Tr}\ln\big[i\partial\!\!\!/ - g_2\frac{\tau^a}{2}W\!\!\!/^a P_L - q_2\frac{1}{6}B\!\!\!/_2 P_L - q_2(\frac{1}{6} + \frac{\tau^3}{2})B\!\!\!/_2 P_R\big]\\
&\quad+iS_{\text{TC2-induced eff}}[U, W, B_2]\Big] \ ,
\end{aligned} \tag{135}$$

Ignoring term $iS_{\text{TC2-induced eff}}[U, W, B_2]$ in above expression, we get expression for $\mathcal{N}[W^a_\mu, B_\mu]$. In above result, if we denote the Z' involved part be $\int\mathcal{D}Z'_\mu\, e^{iS_{Z'}[Z', U, W^a, B]}$, then we will find that Z' field in $S_{Z'}[Z', U, W^a, B]$ is not correctly normalized, since the coefficient in front of kinetic term is not standard $-1/4$. We now introduce normalized fields $Z'_{R,\mu}$ as

$$
\begin{aligned}
Z'_\mu &= \frac{1}{c'} Z'_{R,\mu} && (136)\\
c^{\prime,2} &= 1 + \mathcal{K}\frac{3g_1^2}{2}(\cot^2\theta' + \tan^2\theta') + \frac{3g_1^2}{4}[\mathcal{K}_2^{\mathrm{TC1},\Sigma\neq 0}(\cot\theta' + \tan\theta')^2 && (137)\\
&\quad + \mathcal{K}_{13}^{\mathrm{TC1},\Sigma\neq 0}(\cot\theta' - \tan\theta')^2] - \frac{11}{9}g_1^2\mathcal{K}\tan^2\theta' - \frac{2}{9}(L_{10}^{\mathrm{1D}} + 11H_1^{\mathrm{1D}})g_1^2\tan^2\theta'
\end{aligned}
$$

then

$$
S_{Z'}[Z', U, W^a, B] = \int d^4x\ [\frac{1}{2}Z'_{R,\mu}(x)D_Z^{-1,\mu\nu}Z'_{R,\nu}(x) + Z_R^{\prime,\mu}J_{Z,\mu} + Z_R^2 Z'_{R,\mu}J_{3Z}^\mu + g_{4Z}\frac{g_1^4}{c^{\prime 4}}Z_R^{\prime,4}] \tag{138}
$$

with

$$
\begin{aligned}
D_Z^{-1,\mu\nu} &= g^{\mu\nu}(\partial^2 + M_{Z'}^2) - (1+\lambda_Z)\partial^\mu\partial^\nu + \Delta_Z^{\mu\nu}(X) && (139)\\
M_{Z'}^2 &= \frac{3g_1^2(F_0^{\mathrm{TC1}})^2}{8c^{\prime 2}}(\cot\theta' + \tan\theta')^2 + \frac{g_1^2(F_0^{\mathrm{TC2}})^2}{4c^{\prime 2}}\tan^2\theta' && (140)\\
\lambda_Z &= -\mathcal{K}_1^{\mathrm{TC1},\Sigma\neq 0}\frac{3g_1^2}{8c^{\prime 2}}(\cot\theta' + \tan\theta')^2 && (141)\\
\Delta_Z^{\mu\nu}(X) &= -g_1^2\frac{\tan^2\theta'}{c^{\prime 2}}\{(2L_1^{\mathrm{1D}} + L_3^{\mathrm{1D}})g^{\mu\nu}(\mathrm{tr}X^2) + (2L_1^{\mathrm{1D}} + L_3^{\mathrm{1D}})\mathrm{tr}(X_\mu\tau^3)\mathrm{tr}(X_\nu\tau 3)\\
&\quad + 2L_2^{\mathrm{1D}}\mathrm{tr}(X_\mu X_\nu) + L_2^{\mathrm{1D}}[g^{\mu\nu}\mathrm{tr}(X_\nu\tau^3)\mathrm{tr}(X^\nu\tau^3) + \mathrm{tr}(X_\mu\tau^3)\mathrm{tr}(X_\nu\tau^3)]\} && (142)
\end{aligned}
$$

and

$$
\begin{aligned}
J_Z^\mu &= J_{Z0}^\mu + \frac{g_1^2\gamma}{c'}\partial^\nu B_{\mu\nu} + \tilde{J}_Z^\mu && (143)\\
J_{Z0}^\mu &= -ig_1\frac{(F_0^{\mathrm{TC2}})^2}{4c'}\tan\theta'\mathrm{tr}(\tau^3 X^\mu) && (144)\\
\gamma &= -(3\mathcal{K} + \mathcal{K}_{13}^{\mathrm{TC1},\Sigma\neq 0})\frac{1}{2}(\cot\theta' - \tan\theta') - (11\mathcal{K} + 2L_{10}^{\mathrm{1D}} + 22H_1^{\mathrm{1D}})\frac{1}{9}\tan\theta' && (145)\\
\tilde{J}_{Z,\mu} &= \frac{2i}{c'}(L_1^{\mathrm{1D}} + \frac{L_3^{\mathrm{1D}}}{2})g_1\tan\theta'\mathrm{tr}(X_\mu\tau^3)(\mathrm{tr}X^2) + \frac{2i}{c'}L_2^{\mathrm{1D}}g_1\tan\theta'\mathrm{tr}(X_\nu\tau^3)(\mathrm{tr}X_\mu X_\nu)\\
&\quad + \frac{1}{2c'}g_1 L_9^{\mathrm{1D}}\tan\theta'\mathrm{tr}[(\overline{W}_{\mu\nu}\tau^3 - \tau^3\overline{W}_{\mu\nu})X^\nu] + \frac{i}{2c'}L_9^{\mathrm{1D}}g_1\tan\theta'\partial^\nu[\mathrm{tr}\tau^3(X_\mu X_\nu - X_\nu X_\mu)]\\
&\quad - \frac{1}{c'}L_{10}^{\mathrm{1D}}g_1\tan\theta'\partial^\nu\mathrm{tr}(\tau^3\overline{W}_{\mu\nu})\\
g_{4Z} &= (\mathcal{K}_3^{\mathrm{TC1},\Sigma\neq 0} + \mathcal{K}_4^{\mathrm{TC1},\Sigma\neq 0})\frac{3}{256}(\cot\theta' + \tan\theta')^4 + (L_1^{\mathrm{1D}} + L_2^{\mathrm{1D}} + \frac{L_3^{\mathrm{1D}}}{2})\frac{1}{4}\tan^4\theta' && (146)\\
J_{3Z}^\mu &= -\frac{i}{c^{\prime 3}}(L_1^{\mathrm{1D}} + L_2^{\mathrm{1D}} + \frac{L_3^{\mathrm{1D}}}{2})g_1^3\tan^3\theta'\mathrm{tr}(X_\mu\tau^3) && (147)
\end{aligned}
$$

We denote the result action after the integration over Z' as

$$
\int \mathcal{D}Z'_\mu\ e^{iS_{Z'}[Z',U,W^a,B]} = e^{i\bar{S}_{Z'}[U,W^a,B]} \tag{148}
$$

We can use loop expansion to calculate above integration

$$\bar{S}_{Z'}[U,W^a,B]=S_{Z'}[Z'_c,U,W^a,B]+\text{loop terms} \tag{149}$$

with classical field Z'_c satisfy

$$\frac{\partial}{\partial Z'_{c,\mu}(x)}\Big[S_{Z'}[Z'_c,U,W^a,B]+\text{loop terms}\Big]=0 \tag{150}$$

With (138), the solution is

$$Z'^{\mu}_c(x)=-D^{\mu\nu}_Z J_{Z,\nu}(x)+O(p^3)+\text{loop terms} \tag{151}$$

then

$$\begin{aligned}\bar{S}_{Z'}[U,W^a,B] \;=\; &\int d^4x\,[-\frac{1}{2}J_{Z,\mu}D^{\mu\nu}_Z J_{Z,\nu}-J_{3Z,\mu'}(D^{\mu'\nu'}_Z J_{Z,\nu'})(D^{\mu\nu}_Z J_{Z,\nu})^2+g_{4Z}\frac{g_1^4}{c'^4}(D^{\mu\nu}_Z J_{Z,\nu})^4]\\ &+\text{loop terms}\end{aligned} \tag{152}$$

where

$$D^{-1,\mu\nu}_Z D_{Z,\nu\lambda}=D^{\mu\nu}_Z D^{-1}_{Z,\nu\lambda}=g^{\mu}_{\lambda} \tag{153}$$

and it is not difficult to show that if we are accurate up to order of p^4, then p order Z'_c solution is enough, all contributions from p^3 order Z'_c are at least belong to order of p^6.

With these results, (135) become

$$\begin{aligned}&\mathcal{N}[W^a_\mu,B_\mu]\exp\Big(iS_{\text{eff}}[U,W^a_\mu,B_\mu]\Big) \qquad\qquad (154)\\ &=\exp\bigg[i\bar{S}_{Z'}[U,W^a,B]+iS^{(4c,B)}_{\text{TC1}}[B]+i\int d^4x\Big[-\frac{1}{4}W^a_{\mu\nu}W^{a\mu\nu}-\frac{1}{4}B_{\mu\nu}B^{\mu\nu}-\mathcal{K}\frac{3g_1^2}{4}B_{\mu\nu}B^{\mu\nu}\\ &\quad+\frac{1}{2}\mathcal{K}(\frac{11}{18}g_1^2B_{\mu\nu}B^{\mu\nu}+\frac{1}{2}g_2^2W^a_{\mu\nu}W^{a,\mu\nu})-\frac{(F_0^{\text{TC2}})^2}{4}\text{tr}(X_\mu X^\mu)+(L_1^{\text{1D}}+\frac{L_3^{\text{1D}}}{2})[\text{tr}(X_\mu X^\mu)]^2\\ &\quad+L_2^{\text{1D}}[\text{tr}(X_\mu X_\nu)]^2-iL_9^{\text{1D}}\text{tr}(\overline{W}_{\mu\nu}X^\mu X^\nu)-i\frac{L_9^{\text{1D}}}{2}g_1B_{\mu\nu}\text{tr}(\tau^3X^\mu X^\nu)+\frac{L_{10}^{\text{1D}}}{2}g_1B_{\mu\nu}\text{tr}(\tau^3\overline{W}^{\mu\nu})\\ &\quad+\frac{1}{18}(L_{10}^{\text{1D}}+11H_1^{\text{1D}})g_1^2B_{\mu\nu}B^{\mu\nu}+H_1^{\text{1D}}\text{tr}(\overline{W}_{\mu\nu}\overline{W}^{\mu\nu})\Big]\bigg]\;,\end{aligned}$$

Ignoring term with coefficients F_0^{TC2}, L_i^{1D} and H_1^{1D} in above expression, we get expression for $\mathcal{N}[W^a_\mu,B_\mu]$, with it we finally obtain $S_{\text{eff}}[U,W^a_\mu,B_\mu]$

$$\begin{aligned}S_{\text{eff}}[U,W^a_\mu,B_\mu] \;=\; &\int d^4x\,\Big[-\frac{(F_0^{\text{TC2}})^2}{4}\text{tr}(X_\mu X^\mu)+(L_1^{\text{1D}}+\frac{L_3^{\text{1D}}}{2})[\text{tr}(X_\mu X^\mu)]^2+L_2^{\text{1D}}[\text{tr}(X_\mu X_\nu)]^2\\ &-iL_9^{\text{1D}}\text{tr}(\overline{W}_{\mu\nu}X^\mu X^\nu)-i\frac{L_9^{\text{1D}}}{2}g_1B_{\mu\nu}\text{tr}(\tau^3X^\mu X^\nu)+\frac{L_{10}^{\text{1D}}}{2}g_1B_{\mu\nu}\text{tr}(\tau^3\overline{W}^{\mu\nu})\\ &+\frac{1}{18}(L_{10}^{\text{1D}}+11H_1^{\text{1D}})g_1^2B_{\mu\nu}B^{\mu\nu}+H_1^{\text{1D}}\text{tr}(\overline{W}_{\mu\nu}\overline{W}^{\mu\nu})\Big]+\Delta S_{\text{eff}}[U,W^a_\mu,B_\mu]\end{aligned} \tag{155}$$

i.e. our result EWCL is equal to standard one-doublet technicolor model result plus contributions from Z', we denote this Z' contribution part $\Delta S_{\rm eff}[U,W_\mu^a,B_\mu]$,

$$\Delta S_{\rm eff}[U,W_\mu^a,B_\mu]=\bar{S}_{Z'}[U,W^a,B]-\bar{S}_{Z'}[U,W^a,B]|_{F_0^{\rm TC2}=0,L_i^{\rm 1D}=H_1^{\rm 1D}=0}\ . \tag{156}$$

Correspondingly, EWCL coefficients for topcolor assisted technicolor model can also be divided into two parts

$$f^2=(F_0^{\rm TC2})^2\qquad\beta_1=\Delta\beta_1\qquad\alpha_i=\alpha_i|_{\text{one doulet}}+\Delta\alpha_i\quad i=1,2,\ldots,14 \tag{157}$$

in which $\alpha_i|_{\text{one doulet}}\quad i=1,2,\ldots,14$ are coefficients from one-doublet technicolor model, their values are given in (9) and Table I. $\Delta\beta_1$ and $\Delta\alpha_i\quad i=1,2,\ldots,14$ are contributions from Z' and ordinary quarks, since we do not consider ordinary quarks in this work, so in the next part of this paper, we calculate Z' contributions.

With help of (152), (139) and (143)

$$\begin{aligned}\Delta S_{\rm eff}[U,W_\mu^a,B_\mu]\ =\ &\int d^4x\ [-\frac{1}{2}J_{Z0,\mu}D_Z^{\mu\nu}J_{Z0,\nu}-\frac{1}{M_{Z'}^2}J_{Z0,\mu}(\tilde{J}_Z^\mu+\frac{g_1^2\gamma}{c'}\partial_\nu B^{\mu\nu})\\&-\frac{1}{M_{Z'}^6}J_{3Z,\mu}J_{Z0}^\mu J_{Z0}^2+\frac{g_{4Z}g_1^4}{c'^4M_{Z'}^8}J_{Z0}^4]\end{aligned} \tag{158}$$

With help of following algebra relations,

$$\begin{aligned}&\partial_\mu{\rm tr}[\tau^3X^\mu]=0\\&{\rm tr}[\tau^3(\partial_\mu X_\nu-\partial_\nu X_\mu)]=-2{\rm tr}(\tau^3X_\mu X_\nu)+i{\rm tr}(\tau^3\overline{W}_{\mu\nu})-ig_1B_{\mu\nu}\\&{\rm tr}(\tau^3X_\mu X_\nu){\rm tr}(\tau^3X^\mu X^\nu)\\&=[{\rm tr}(X_\mu X_\nu)]^2-[{\rm tr}(X_\mu X^\mu)]^2-{\rm tr}(X_\mu X_\nu){\rm tr}(\tau^3X^\mu){\rm tr}(\tau^3X^\nu)+{\rm tr}(X_\mu X^\mu)[{\rm tr}(\tau^3X_\nu)]^2\\&{\rm tr}(TA){\rm tr}(TBC)+{\rm tr}(TB){\rm tr}(TCA)+{\rm tr}(TC){\rm tr}(TAB)=2{\rm tr}(ABC)\end{aligned} \tag{159}$$

where ${\rm tr}A={\rm tr}B={\rm tr}C=0$ and $T^2=1$. We can show (158) leads to the form of standard EWCL, further combined with (155) and (144), we can read out p^2 coefficient

$$\beta_1=\frac{g_1^2(F_0^{\rm TC2})^2}{8c'^2M_{Z'}^2}\tan^2\theta'=\frac{(F_0^{\rm TC2})^2}{3(F_0^{\rm TC1})^2(\cot^2\theta'+1)^2+2(F_0^{\rm TC2})^2} \tag{160}$$

which imply a positive β_1 which is further bounded above. With the fact that $f=F_0^{\rm TC2}=250$GeV and original model requirement $F_0^{\rm TC1}=1$TeV, we find

$$2\beta_1=\frac{1}{24(\cot^2\theta'+1)^2+1}\ . \tag{161}$$

Combine with $\alpha T = 2\beta_1$ given in Ref.[2], we obtain result that topcolor-assisted technicolor model produce positive and bounded above T parameter! The upper limit of β_1 is $1/50$ which corresponds to upper limit of T parameter $1/(25\alpha) \sim 5.1$. From (160), we know β_1 coefficient is uniquely determined by parameter θ', therefore instead of using θ' as the input parameter of the theory, we can further use β_1 or $T = 2\beta_1/\alpha$ as the parameter of the theory.

The p^4 order coefficients can be read out from derived EWCL (155) and (158), we list down the results as following,

$$
\begin{aligned}
&\alpha_1 = (1-2\beta_1)L_{10}^{\mathrm{1D}} + \frac{(F_0^{\mathrm{TC2}})^2}{2M_{Z'}^2}\beta_1 - 2\gamma\beta_1\cot\theta' \\
&\alpha_2 = -\frac{1}{2}(1-2\beta_1)L_9^{\mathrm{1D}} - \frac{(F_0^{\mathrm{TC2}})^2}{2M_{Z'}^2}\beta_1 - 2\gamma\beta_1\cot\theta' \\
&\alpha_3 = -\frac{1}{2}(1-2\beta_1)L_9^{\mathrm{1D}} \qquad\qquad \alpha_4 = L_2^{\mathrm{1D}} + \frac{(F_0^{\mathrm{TC2}})^2}{2M_{Z'}^2}\beta_1 + 2\beta_1 L_9^{\mathrm{1D}} \\
&\alpha_5 = L_1^{\mathrm{1D}} + \frac{L_3^{\mathrm{1D}}}{2} - \frac{(F_0^{\mathrm{TC2}})^2}{2M_{Z'}^2}\beta_1 - 2\beta_1 L_9^{\mathrm{1D}} \\
&\alpha_6 = -\frac{(F_0^{\mathrm{TC2}})^2}{2M_{Z'}^2}\beta_1 + 4\beta_1^2 L_2^{\mathrm{1D}} - 4\beta_1(L_2^{\mathrm{1D}} + \frac{L_9^{\mathrm{1D}}}{2}) \\
&\alpha_7 = \beta_1\frac{(F_0^{\mathrm{TC2}})^2}{2M_{Z'}^2} + 2(\beta_1^2 - \beta_1)(2L_1^{\mathrm{1D}} + L_3^{\mathrm{1D}}) + 2\beta_1 L_9^{\mathrm{1D}} \\
&\alpha_8 = -\beta_1\frac{(F_0^{\mathrm{TC2}})^2}{2M_{Z'}^2} + \beta_1 L_{10}^{\mathrm{1D}} \\
&\alpha_9 = -\beta_1\frac{(F_0^{\mathrm{TC2}})^2}{2M_{Z'}^2} + 2\beta_1(-L_9^{\mathrm{1D}} + L_{10}^{\mathrm{1D}}) \\
&\alpha_{10} = (4\beta_1^2 - 8\beta_1^3)(L_1^{\mathrm{1D}} + L_2^{\mathrm{1D}} + \frac{L_3^{\mathrm{1D}}}{2}) + 16\beta_1^4 g_{4Z}\cot^4\theta' \\
&\alpha_{11} = \alpha_{12} = \alpha_{13} = \alpha_{14} = 0
\end{aligned}
\tag{162}
$$

Several features of this result are:

1. Except the part of one-doublet technicolor model result, all corrections from Z' particle are at least proportional to β_1 which vanish if the mixing disappear by $\theta' = 0$.

2. Since $L_{10}^{\mathrm{1D}} < 0$, therefore (162) tells us α_8 is negative and then $U = -16\pi\alpha_8$ is always positive in this model.

3. Except α_1, α_2 and α_{10}, all other coefficients are determined by one-doublet technicolor model coefficients given in Table VI and two other parameter β_1 and $F_0^{\mathrm{TC2}}/M_{Z'}$.

4. α_{10} further depend on parameter g_{4Z} which from (146) further depend on $\mathcal{K}_3^{\mathrm{TC1},\Sigma\neq 0}+\mathcal{K}_4^{\mathrm{TC1},\Sigma\neq 0}$ which are already given by (101) and Table III.

5. α_1 and α_2 further depend on γ which from (145) further rely on an extra parameter $\mathcal{K}$. We can combine (160) and (137) together to fix $\mathcal{K}$,

$$\begin{aligned}\frac{(F_0^{\mathrm{TC2}})^2}{8\beta_1 M_{Z'}^2}\tan^2\theta' &= \frac{1}{g_1^2}+\mathcal{K}(\frac{3}{2}\cot^2\theta'+\frac{5}{18}\tan^2\theta')+\frac{3}{4}[\mathcal{K}_2^{\mathrm{TC1},\Sigma\neq 0}(\cot\theta'+\tan\theta')^2\\ &\quad +\mathcal{K}_{13}^{\mathrm{TC1},\Sigma\neq 0}(\cot\theta'-\tan\theta')^2]-\frac{2}{9}(L_{10}^{\mathrm{1D}}+11H_1^{\mathrm{1D}})\tan^2\theta'\end{aligned}$$

   Once $\mathcal{K}$ is fixed, with help of (92), we can determine the ratio of infrared cutoff κ and ultraviolet cutoff Λ, in Fig.1(a), we draw the κ/Λ as function of T and $M_{Z'}$, we find our calculation do produce very large hierarchy and we further find not all T and $M_{Z'}$ region is available if we consider the natural criteria $\Lambda > \kappa$. This criteria leads constraints that as long as Z' mass become large, the allowed range for T parameter become smaller and smaller approaching to zero, for example, $T < 0.37$ for $M_{Z'} = 0.5$TeV, $T < 0.0223$ for $M_{Z'} = 1$TeV and $T < 0.004$ for $M_{Z'} = 2$TeV. In Fig.1(b), we draw Z' mass as function of T parameter and κ/Λ. The line of $\kappa/\Lambda = 1$ give the upper bound of Z' mass. The upper bound of Z' mass depend on value of T parameter, the smaller the T, the larger the upper bound of $M_{Z'}$.

6. For fixed $M_{Z'}$, there exists a special θ' value which maximizes α_1. The parameter, $S = -16\pi\alpha_1$, is of special importance in new physics search, in Fig.2, we draw a graph of minimal S parameter with different T parameter. We see that if the Z' mass is low enough, say $M_{Z'} < 0.441$ TeV or $T > 0.176$, S will become negative.

Since we already know $F_0^{\mathrm{TC2}} = 250$GeV, therefore all EWCL coefficients depend on two physical parameters β_1 and $M_{Z'}$. Combined with $\alpha T = 2\beta_1$, we can use the present experimental result for the T parameter to fix β_1. In Fig.3, we draw graphs for the S and U parameters in terms of the T parameter. We take three typical Z' masses $M_{Z'} = 0.5,\ 1,\ 2$ TeV for references. In Fig.4, we draw graphs for all p^4 order nonzero coefficients in terms of the T parameter. Where for α_3 and α_{10}, we only draw one line for each of them, since they are independent of the Z' mass.

FIG. 1: (a). The ratio of infrared cutoff and ultraviolet cutoff κ/Λ as function of T parameter and Z' mass in unit of TeV. (b). Z' mass in unit of TeV as function of T parameter and κ/Λ.



(a) κ/Λ

(b) $M_{Z'}$

IV. CONCLUSION

In this paper, we have set up a formulation to perform the dynamical computation of the bosonic part of EWCL for the one-doublet and topcolor-assisted technicolor models. The one-doublet technicolor model as the earliest and simplest dynamical symmetry breaking model are taken as the trial model to test our formulation. We find our formulation recovers standard scaling-up results. The topcolor-assisted technicolor model is the main model we handle in this paper. We have computed its TC1 dynamics in detail and verify the dynamical symmetry breaking of the theory, TC1 interaction will induce effective interactions among colorons and Z' which are characterized by a divergent constant $\mathcal{K}$, a dimensional constant $F_0^{\rm TC1}$ and a series of dimensionless QCD constants $L_1, L_3, L_9, L_{10}, H_1$. For TC2 dynamics, it will induce effective interaction for Z', electroweak gauge fields and their goldstone bosons. Due to its similarity with QCD, we use Gasser-Leutwyler prescription to describe its low energy effects in terms of low energy effective Lagrangian with a divergent constant $\mathcal{K}$, dimensional constant $F_0^{\rm TC2}$ and a series of dimensionless constants $L_1^{\rm 1D}, L_2^{\rm 1D}, L_3^{\rm 1D}, L_9^{\rm 1D}, L_{10}^{\rm 1D}, H_1^{\rm 1D}$ the same as those in one-doublet technicolor model. Due to

FIG. 2: The dashed line is the minimal S parameter in topcolor assisted technicolor model for different T. The solid lines are the isolines for different choices of Z' mass in units of TeV.



dynamics similarities between TC2 and QCD, TC2 interaction make a direct contributions to EWCL coefficients a part which is the same as that of one-doublet technicolor model. Further corrections are from effective interactions among colorons, Z' and ordinary quarks induced by TC1 and TC2 interactions. We have shown that colorons make no contributions to EWCL coefficients within the approximations we have made in this paper, while ordinary quark are ignored in this paper for future investigations. In fact, for some special EWCL coefficients, such as $S = -16\pi\alpha_1$, $\alpha T = \beta_1$ and $U = -16\pi\alpha_8$ parameters, general fermion contributions to them are already calculated [23], S, T, U and triple-gauge-vertices from a heavy non-degenerate fermion doublet has been estimated in ref.[9, 24]. One can based on these general results to estimate possible contributions to some of EWCL coefficients. For topcolor-assisted technicolor model in this paper, the main work is to estimate the effects of Z' particle. Our computation shows that contributions from Z' particle are at least proportional to β_1 and then vanish if β_1 is zero. One typical feature of the model is the positivity and bounded above of β_1 parameter which means the T parameter must vary in the range $0 \sim 1/(25\alpha)$ and the positive U parameter. If we consider the natural criteria $\Lambda > \kappa$ which will further constraints the allowed range for T parameter approaching to zero as long as Z' mass become large, for example, $T < 0.37$ for $M_{Z'} = 0.5$TeV, $T < 0.0223$ for $M_{Z'} = 1$TeV and $T < 0.004$ for $M_{Z'} = 2$TeV. For S parameter, it can be either positive and negative

FIG. 3: The S and U parameters for topcolor assisted technicolor model. $F_0^{\rm TC2} = 250$ GeV, the T parameter and $M_{Z'} = \{0.5,\ 1,\ 2\}$ TeV are as input parameters of the model.



(a) S

(b) U

depending on the Z' mass is large or small. As long as $M_{Z'} < 0.441$ TeV or $T > 0.176$, we may find negative S. There exist a upper bound for the mass of Z' which is depend on value of T parameter, the smaller the T, the larger the upper bound of $M_{Z'}$. Except $U(1)_Y$ coupling g_1 and coefficients determined in one-doublet technicolor model and QCD, all EWCL coefficients rely on experimental T parameter and coloron mass $M_{Z'}$. We have taken typical values of $M_{Z'}$ and vary T parameter to estimate all EWCL coefficients up to order of p^4. Further works on matter part of EWCL and computing EWCL coefficients for other dynamical symmetry breaking new physics models are in progress and will be reported elsewhere.

Acknowledgments

We would like to thank Y. P. Kuang, H. J. He and J. K. Parry for helpful discussions. This work was supported by National Science Foundation of China (NSFC) under Grant No. 10435040 and Specialized Research Fund for the Doctoral Program of High Education of China.

FIG. 4: Nonzero EWCL coefficients $\alpha_i(i = 1, 2, \ldots, 10)$ for the topcolor-assisted technicolor model up to order of $\mathcal{O}(p^4)$. $F_0^{\mathrm{TC2}} = 250$ GeV, the T parameter and $M_{Z'} = \{0.5,\ 1,\ 2\}$ TeV are as input parameters of the model.



(a) α_1 (b) α_2

(c) α_3 (d) α_4

APPENDIX A: TWO EQUIVALENT EWCL FORMALISMS

The electroweak chiral Lagrangian is constructed using a dimensionless unitary unimodular 2×2 matrix field $U(x)$, In Ref.[2], the electroweak chiral Lagrangian has been constructed

FIG. 4:



(a) α_5 (b) α_6



(c) α_7 (d) α_8

with the building blocks which are $SU(2)_L$ covariant and $U(1)_Y$ invariant as:

$$T \equiv U\tau^3U^\dagger\ , \qquad V_\mu \equiv (D_\mu U)U^\dagger\ , \qquad g_1B_{\mu\nu}\ , \qquad g_2W_{\mu\nu} \equiv g_2\frac{\tau^a}{2}W^a_{\mu\nu}\ . \tag{A1}$$

Alternatively, we reformulate the electroweak chiral Lagrangian equivalently with $SU(2)_L$ invariant and $U(1)_Y$ covariant building blocks as:

$$\tau^3\ , \qquad X_\mu \equiv U^\dagger(D_\mu U)\ , \qquad g_1B_{\mu\nu}\ , \qquad \overline{W}_{\mu\nu} \equiv U^\dagger g_2W_{\mu\nu}U\ , \tag{A2}$$

FIG. 4:



(a) α_9

(b) α_{10}

among which, τ^3 and $g_1 B_{\mu\nu}$ are both $SU(2)_L$ and $U(1)_Y$ invariant, while X_μ and $\overline{W}_{\mu\nu}$ are bilinearly $U(1)_Y$ covariant. This second formulation is largely used throughout this paper. We list down the corresponding relations of the two formalisms.

APPENDIX B: $S_{\rm TC1}^{(4c,B^A)}[B^A]$, $S_{\rm TC1}^{(4c,B)}[B]$, $S_{\rm TC1}^{(4c,Z')}$, $S_{\rm TC1}^{(4c,B^AZ')}[B^A,Z']$, $S_{\rm TC1}^{(4c,BZ')}[B,Z']$

$$\begin{aligned}
S_{\rm TC1}^{(4c,B^A)}[B^A] \;=\; \int d^4x\bigg[&-\mathcal{K}_1^{\rm TC1,\Sigma\neq0}\frac{g_3^2}{8}(\cot\theta+\tan\theta)^2(\partial_\mu B^{A,\mu})^2\\
&-\mathcal{K}_2^{\rm TC1,\Sigma\neq0}\frac{g_3^2}{8}(\cot\theta+\tan\theta)^2B^A_{a,\mu\nu}B^{A,\mu\nu}_a\\
&+\mathcal{K}_3^{\rm TC1,\Sigma\neq0}[\frac{g_3^4}{192}(\cot\theta+\tan\theta)^4(B^A_\mu B^{A,\mu})^2+\frac{g_3^4}{128}(\cot\theta+\tan\theta)^4(d^{ABC}B^B_\mu B^{C,\mu})^2]\\
&+\mathcal{K}_4^{\rm TC1,\Sigma\neq0}\{\frac{g_3^4}{192}(\cot\theta+\tan\theta)^4(B^A_\mu B^A_\nu)^2\\
&+\frac{g_3^4}{128}(\cot\theta+\tan\theta)^4[(if^{ABC}+d^{ABC})B^B_\mu B^C_\nu]^2\}\\
&-\mathcal{K}_{13}^{\rm TC1,\Sigma\neq0}\frac{g_3^2}{8}(\cot\theta-\tan\theta)^2B^A_{v,\mu\nu}B^{A,\mu\nu}_v\\
&+\mathcal{K}_{14}^{\rm TC1,\Sigma\neq0}\frac{g_3^3}{32}(\cot\theta-\tan\theta)(\cot\theta+\tan\theta)^2B^{A,\mu\nu}_vf^{ABC}B^B_\mu B^C_\nu\bigg]\;, \qquad \text{(B1)}
\end{aligned}$$

TABLE I: The Symmetry Breaking Sector of the Electroweak Chiral Lagrangian

	Formulation I	Formulation II
$\mathcal{L}^{(2)}$	$\frac{1}{4}f^2\mathrm{tr}[(D_\mu U^\dagger)(D^\mu U)] = -\frac{1}{4}f^2\mathrm{tr}(V_\mu V^\mu)$	$-\frac{1}{4}f^2\mathrm{tr}(X_\mu X^\mu)$
$\mathcal{L}^{(2)\prime}$	$\frac{1}{4}\beta_1 f^2[\mathrm{tr}(TV_\mu)]^2$	$\frac{1}{4}\beta_1 f^2[\mathrm{tr}(\tau^3 X_\mu)]^2$
$\mathcal{L}_1$	$\frac{1}{2}\alpha_1 g_2 g_1 B_{\mu\nu}\mathrm{tr}(TW^{\mu\nu})$	$\frac{1}{2}\alpha_1 g_1 B_{\mu\nu}\mathrm{tr}(\tau^3\overline{W}^{\mu\nu})$
$\mathcal{L}_2$	$\frac{1}{2}i\alpha_2 g_1 B_{\mu\nu}\mathrm{tr}(T[V^\mu, V^\nu])$	$i\alpha_2 g_1 B_{\mu\nu}\mathrm{tr}(\tau^3 X^\mu X^\nu)$
$\mathcal{L}_3$	$i\alpha_3 g_2\mathrm{tr}(W_{\mu\nu}[V^\mu, V^\nu])$	$2i\alpha_3\mathrm{tr}(\overline{W}_{\mu\nu}X^\mu X^\nu)$
$\mathcal{L}_4$	$\alpha_4[\mathrm{tr}(V_\mu V_\nu)]^2$	$\alpha_4[\mathrm{tr}(X_\mu X_\nu)]^2$
$\mathcal{L}_5$	$\alpha_5[\mathrm{tr}(V_\mu V^\mu)]^2$	$\alpha_5[\mathrm{tr}(X_\mu X^\mu)]^2$
$\mathcal{L}_6$	$\alpha_6\mathrm{tr}(V_\mu V_\nu)\mathrm{tr}(TV^\mu)\mathrm{tr}(TV^\nu)$	$\alpha_6\mathrm{tr}(X_\mu X_\nu)\mathrm{tr}(\tau^3 X^\mu)\mathrm{tr}(\tau^3 X^\nu)$
$\mathcal{L}_7$	$\alpha_7\mathrm{tr}(V_\mu V^\mu)\mathrm{tr}(TV_\nu)\mathrm{tr}(TV^\nu)$	$\alpha_7\mathrm{tr}(X_\mu X^\mu)\mathrm{tr}(\tau^3 X_\nu)\mathrm{tr}(\tau^3 X^\nu)$
$\mathcal{L}_8$	$\frac{1}{4}\alpha_8 g_2^2[\mathrm{tr}(TW_{\mu\nu})]^2$	$\frac{1}{4}\alpha_8[\mathrm{tr}(\tau^3\overline{W}_{\mu\nu})]^2$
$\mathcal{L}_9$	$\frac{1}{2}i\alpha_9 g_2\mathrm{tr}(TW_{\mu\nu})\mathrm{tr}(T[V^\mu, V^\nu])$	$i\alpha_9\mathrm{tr}(\tau^3\overline{W}_{\mu\nu})\mathrm{tr}(\tau^3 X^\mu X^\nu)$
$\mathcal{L}_{10}$	$\frac{1}{2}\alpha_{10}[\mathrm{tr}(TV_\mu)\mathrm{tr}(TV_\nu)]^2$	$\frac{1}{2}\alpha_{10}[\mathrm{tr}(\tau^3 X_\mu)\mathrm{tr}(\tau^3 X_\nu)]^2$
$\mathcal{L}_{11}$	$\alpha_{11}g_2\epsilon^{\mu\nu\rho\lambda}\mathrm{tr}(TV_\mu)\mathrm{tr}(V_\nu W_{\rho\lambda})$	$\alpha_{11}\epsilon^{\mu\nu\rho\lambda}\mathrm{tr}(\tau^3 X_\mu)\mathrm{tr}(X_\nu\overline{W}_{\rho\lambda})$
$\mathcal{L}_{12}$	$\alpha_{12}g_2\mathrm{tr}(TV_\mu)\mathrm{tr}(V_\nu W^{\mu\nu})$	$\alpha_{12}\mathrm{tr}(\tau^3 X_\mu)\mathrm{tr}(X_\nu\overline{W}^{\mu\nu})$
$\mathcal{L}_{13}$	$\alpha_{13}g_2 g_1\epsilon^{\mu\nu\rho\sigma}B_{\mu\nu}\mathrm{tr}(TW_{\rho\sigma})$	$\alpha_{13}\epsilon^{\mu\nu\rho\sigma}g_1 B_{\mu\nu}\mathrm{tr}(\tau^3\overline{W}_{\rho\sigma})$
$\mathcal{L}_{14}$	$\alpha_{14}g_2^2\epsilon^{\mu\nu\rho\sigma}\mathrm{tr}(TW_{\mu\nu})\mathrm{tr}(TW_{\rho\sigma})$	$\alpha_{14}\epsilon^{\mu\nu\rho\sigma}\mathrm{tr}(\tau^3\overline{W}_{\mu\nu})\mathrm{tr}(\tau^3\overline{W}_{\rho\sigma})$

$$S_{\mathrm{TC1}}^{(4c,B)}[B] = \int d^4x\Big[-\mathcal{K}_{13}^{\mathrm{TC1},\Sigma\neq 0}\frac{3g_1^2}{4}B_{\mu\nu}B^{\mu\nu}\Big]\,, \tag{B2}$$

$$\begin{aligned} S_{\mathrm{TC1}}^{(4c,Z')} = \int d^4x\,\Big[& -\mathcal{K}_1^{\mathrm{TC1},\Sigma\neq 0}\frac{3g_1^2}{16}(\cot\theta'+\tan\theta')^2(\partial_\mu Z'^{,\mu})^2 \\ & +(\mathcal{K}_3^{\mathrm{TC1},\Sigma\neq 0}+\mathcal{K}_4^{\mathrm{TC1},\Sigma\neq 0})\frac{3g_1^4}{256}(\cot\theta'+\tan\theta')^4(Z'_\mu Z'^{,\mu})^2 \\ & -\frac{3g_1^2}{16}[\mathcal{K}_2^{\mathrm{TC1},\Sigma\neq 0}(\cot\theta'+\tan\theta')^2+\mathcal{K}_{13}^{\mathrm{TC1},\Sigma\neq 0}(\cot\theta'-\tan\theta')^2]Z'_{\mu\nu}Z'^{,\mu\nu}\Big]\,, \end{aligned} \tag{B3}$$

$$\begin{aligned}S^{(4c,B^AZ')}_{\rm TC1}[B^A,Z'] \;=\; \int d^4x\;\Big[&\mathcal{K}_3^{\rm TC1,\Sigma\neq 0}[\frac{g_1^2g_3^2}{64}(\cot\theta+\tan\theta)^2(\cot\theta'+\tan\theta')^2 B^A_\mu B^{A,\mu} Z'_\nu Z'^{,\nu}\\ &+\frac{g_1^2g_3^2}{32}(\cot\theta+\tan\theta)^2(\cot\theta'+\tan\theta')^2(B^A_\mu Z'^{,\mu})^2\\ &+\frac{g_1g_3^3}{32}(\cot\theta+\tan\theta)^3(\cot\theta'+\tan\theta')d^{ABC}B^B_\mu B^{C,\mu}B^A_\mu Z'^{,\mu}]\\ &+\mathcal{K}_4^{\rm TC1,\Sigma\neq 0}\{\frac{g_1^2g_3^2}{256}(\cot\theta+\tan\theta)^2(\cot\theta'+\tan\theta')^2(B^A_\mu Z'^{,\mu})^2\\ &+\frac{g_1^2g_3^2}{64}(\cot\theta+\tan\theta)^2(\cot\theta'+\tan\theta')^2[B^A_\mu B^{A,\mu}Z'_\nu Z'^{,\nu}+(B^A_\mu Z'^{,\mu})^2]\\ &+\frac{g_1g_3^3}{32}(\cot\theta+\tan\theta)^3(\cot\theta'+\tan\theta')d^{ABC}B^B_\mu B^C_\nu B^{A,\mu}Z'^{,\nu}\}\Big]\,, \end{aligned} \tag{B4}$$

$$S^{(4c,BZ')}_{\rm TC1}[B,Z'] \;=\; \int d^4x\Big[-\mathcal{K}_{13}^{\rm TC1,\Sigma\neq 0}\frac{g_1^2}{4}(\cot\theta'-\tan\theta')B_{\mu\nu}Z'^{,\mu\nu}\Big]\,, \tag{B5}$$

with $B^A_{v,\mu\nu}=\partial_\mu B^A_\nu-\partial_\nu B^A_\mu-\frac{g_3}{2}(\cot\theta-\tan\theta)f^{ABC}B^B_\mu B^C_\nu$.

APPENDIX C: $S^0_{\rm coloron}[B^A,Z']$ AND $S^{\rm int}_{\rm coloron}[B^A,Z']$

$$\begin{aligned}&S^0_{\rm coloron}[B^A,Z']\\ &=\int d^4x\Big[\Big(g^{\mu\nu}\{\frac{(F_0^{\rm TC1})^2}{8}g_3^2(\cot\theta+\tan\theta)^2+[\mathcal{K}_3^{\rm TC1,\Sigma\neq 0}\frac{g_1^2g_3^2}{64}(\cot\theta+\tan\theta)^2(\cot\theta'+\tan\theta')^2\\ &\quad+\mathcal{K}_4^{\rm TC1,\Sigma\neq 0}\frac{g_1^2g_3^2}{64}(\cot\theta+\tan\theta)^2(\cot\theta'+\tan\theta')^2]Z'_\lambda Z'^{,\lambda}\}\\ &\quad+[\mathcal{K}_3^{\rm TC1,\Sigma\neq 0}\frac{g_1^2g_3^2}{32}(\cot\theta+\tan\theta)^2(\cot\theta'+\tan\theta')^2\\ &\quad+\mathcal{K}_4^{\rm TC1,\Sigma\neq 0}\frac{5g_1^2g_3^2}{256}(\cot\theta+\tan\theta)^2(\cot\theta'+\tan\theta')^2]Z'^{,\mu}Z'^{,\nu}\Big)B^A_\mu B_{A,\nu}\\ &\quad-\mathcal{K}_1^{\rm TC1,\Sigma\neq 0}\frac{g_3^2}{8}(\cot\theta+\tan\theta)^2(\partial_\mu B^{A,\mu})^2\\ &\quad[-\mathcal{K}_2^{\rm TC1,\Sigma\neq 0}\frac{g_3^2}{8}(\cot\theta+\tan\theta)^2-\mathcal{K}_{13}^{\rm TC1,\Sigma\neq 0}\frac{g_3^2}{8}(\cot\theta-\tan\theta)^2\\ &\quad-\frac{1}{4}-\frac{g_3^2}{4}\mathcal{K}(\cot^2\theta+\tan^2\theta)](\partial_\mu B^A_\nu-\partial_\nu B^A_\mu)(\partial^\mu B^{A,\nu}-\partial^\nu B^{A,\mu})\Big]\end{aligned} \tag{C1}$$

and

$$
\begin{aligned}
&S^{\rm int}_{\rm coloron}[B^A,Z'] \qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad \text{(C2)}\\
&=\int d^4x\bigg[-\mathcal{K}_2^{\rm TC1,\Sigma\neq0}\frac{g_3^2}{8}(\cot\theta+\tan\theta)^2[(\partial_\mu B^A_\nu-\partial_\nu B^A_\mu)2g_3(-\cot\theta+\tan\theta)f^{ABC}B^{B,\mu}B^{C,\nu}\\
&\quad+g_3^2(-\cot\theta+\tan\theta)^2f^{ABC}B^B_\mu B^C_\nu f^{AB'C'}B^{B',\mu}B^{C',\nu}]\\
&\quad+\mathcal{K}_3^{\rm TC1,\Sigma\neq0}[\frac{g_3^4}{192}(\cot\theta+\tan\theta)^4(B^A_\mu B^{A,\mu})^2+\frac{g_3^4}{128}(\cot\theta+\tan\theta)^4(d^{ABC}B^B_\mu B^{C,\mu})^2]\\
&\quad+\mathcal{K}_4^{\rm TC1,\Sigma\neq0}\{\frac{g_3^4}{192}(\cot\theta+\tan\theta)^4(B^A_\mu B^A_\nu)^2+\frac{g_3^4}{128}(\cot\theta+\tan\theta)^4[(if^{ABC}+d^{ABC})B^B_\mu B^C_\nu]^2\}\\
&\quad-\mathcal{K}_{13}^{\rm TC1,\Sigma\neq0}\frac{g_3^2}{8}(\cot\theta-\tan\theta)^2[(\partial_\mu B^A_\nu-\partial_\nu B^A_\mu)g_3(-\cot\theta+\tan\theta)f^{ABC}B^{B,\mu}B^{C,\nu}\\
&\quad+\frac{1}{4}g_3^2(-\cot\theta+\tan\theta)^2f^{ABC}B^B_\mu B^C_\nu f^{AB'C'}B^{B',\mu}B^{C',\nu}]+\mathcal{K}_{14}^{\rm TC1,\Sigma\neq0}\frac{g_3^3}{32}(\cot\theta-\tan\theta)(\cot\theta+\tan\theta)^2\\
&\quad\times[\partial^\mu B^{A,\nu}-\partial^\nu B^{A,\mu}+\frac{g_3}{2}(-\cot\theta+\tan\theta)f^{ABC}B^{B,\mu}B^{C,\nu}]f^{ABC}B^B_\mu B^C_\nu\\
&\quad+\mathcal{K}_3^{\rm TC1,\Sigma\neq0}\frac{g_1g_3^3}{32}(\cot\theta+\tan\theta)^3(\cot\theta'+\tan\theta')d^{ABC}B^B_\mu B^{C,\mu}B^A_\nu Z'^{,\nu}\\
&\quad+\mathcal{K}_4^{\rm TC1,\Sigma\neq0}\frac{g_1g_3^3}{32}(\cot\theta+\tan\theta)^3(\cot\theta'+\tan\theta')d^{ABC}B^B_\mu B^C_\nu B^{A,\mu}Z'^{,\nu}\\
&\quad-\frac{1}{4}\cos^2\theta[2g_3(\partial_\mu B^A_\nu-\partial_\nu B^A_\mu)\cot\theta f^{ABC}B^{B,\mu}B^{C,\nu}+g_3^2\cot^2\theta f^{ABC}B^{B,\mu}B^{C,\nu}f^{AB'C'}B^{B'}_\mu B^{C'}_\nu]\\
&\quad-\frac{1}{4}\sin^2\theta[-2g_3(\partial_\mu B^A_\nu-\partial_\nu B^A_\mu)\tan\theta f^{ABC}B^{B,\mu}B^{C,\nu}+g_3^2\tan^2\theta f^{ABC}B^{B,\mu}B^{C,\nu}f^{AB'C'}B^{B'}_\mu B^{C'}_\nu]\\
&\quad-\frac{g_3^2}{4}\mathcal{K}\cot^2\theta[-2g_3(\partial_\mu B^A_\nu-\partial_\nu B^A_\mu)\cot\theta f^{ABC}B^{B,\mu}B^{C,\nu}+g_3^2\cot^2\theta f^{ABC}B^{B,\mu}B^{C,\nu}f^{AB'C'}B^{B'}_\mu B^{C'}_\nu]\\
&\quad-\frac{g_3^2}{4}\mathcal{K}\tan^2\theta[2g_3(\partial_\mu B^A_\nu-\partial_\nu B^A_\mu)\tan\theta f^{ABC}B^{B,\mu}B^{C,\nu}+g_3^2\tan^2\theta f^{ABC}B^{B,\mu}B^{C,\nu}f^{AB'C'}B^{B'}_\mu B^{C'}_\nu]\bigg]\,,
\end{aligned}
$$

---

[1] T.Appelquist and C.Bernard, Phys. Rev. **D22**, 200(1980);
A.Longhitano, Phys. Rev. **D22**, 1166(1980); Nucl. Phys. **B188**,118(1981)

[2] T.Appelquist and G-H. Wu, Phys. Rev. **D48**, 3235(1993); **D51**, 240(1995)

[3] E.Bagan, D.Espriu and J.Manzano, Phys. Rev. **D60**, 114035(1999)

[4] S.Alam, S.Dawson, R.Szalapski, Phys. Rev. **D57**, 1577(1998)

[5] H.-J. He, Y.-P.Kuang, C.-P.Yuan, hep-ph/9704276

[6] A.Dobado and M.T.Urdiales, hep-ph/9502255

[7] A.Nyffeler, A.Schenk, Phys. Rev. **D62**, 113006(2000)

[8] M.J.Herrero and E.R.Morales, hep-ph/9308276

[9] M.E.Peskin and T.Takeuchi, Phys. Rev. Lett. **65**, 964(1990); Phys. Rev. **D46**, 381(1992)

[10] W.-M.Yao, *et al.* (Particle Data Group), J. Phys. **G33**, 1(2006)

[11] Q.Wang, Y.-P.Kuang, M.Xiao and X.-L.Wang, Phys. Rev. **D61**, 054011(2000)

[12] H.Yang, Q.Wang, Y.-P.Kuang and Q.Lu, Phys. Rev. **D66**, 014019(2002)

[13] H.Yang, Q.Wang and Q.Lu, Phys. Lett. **B532**, 240(2002)

[14] Y.-L.Ma, Q.Wang, Phys. Lett. **B560**, 188(2003)

[15] Z.-M.Wang and Q.Wang, Commun. Theor. Phys. 36 ,417(2001)

H.-H.Zhang, W.-B.Yan, J.K.Parry and X.-S.Li, arXiv:0704.1075 [hep-ph]

[16] S.Weinberg, Phys. Rev. **D13**, 974(1976); **D19**, 1277(1979);

[17] C.T.Hill and E.H.Simmons, Phys. Rept. **381**, 235(2003); Erratum-ibid. **390**, 553(2004)

[18] L.Susskind, Phys. Rev. **D20**, 2619(1979);

[19] E.Farhi and L.Susskind, Phys. Rept. **74**, 277(1981);

[20] J.Gasser and H.Leutwyler, Ann. Phys. **158**, 142(1984); Nucl. Phys. **B250**, 465(1985);

[21] C.T.Hill, Phys. Lett. **B345**, 483(1995)

[22] W.-M.Yao, *et al.*, J. Phys. **G33**, 1(2006)

[23] M.J.Dugan and L.Randall, Phys. Lett. **B264**, 154(1991);

H.-H.Zhang, Y.Cao and Q.Wang, hep-ph/0610094

[24] T.Inami, C.S.Lim, B.Takeuchi and M.Tanabashi, Phys. Lett. **B381**, 458(1996)